arXiv:1205.4507v1 [math.AP] 21 May 2012

GLOBAL EXISTENCE OF STRONG SOLUTIONS TO MICROPOLAR EQUATIONS IN CYLINDRICAL DOMAINS

BERNARD NOWAKOWSKI

Abstract. The micropolar equations are a useful generalization of the classical Navier-Stokes model for fluids with micro-structure. We prove the existence of global and strong solutions to these equations in cylindrical domains in $\mathbb{R}^3$. We do not impose any restrictions on the magnitude of the initial and external data but we require that they cannot change in the x_3-direction too fast.

1. Introduction

Introduced in 1966 by A. Eringnen (see [Eri66]), micropolar equations became an important generalization of the classical Navier-Stokes model. These equations take into account that fluid molecules may rotate independently of the fluid rotation. Thus, the standard Navier-Stokes system is complemented with another vector equation which describes the angular momentum of the particles. If we denote the velocity field by v and the microrotation fields by ω, then we see that (v,ω) has six degrees of freedom. Let us clearly emphasize that ω does not represent the rotation field ($\operatorname{rot} v$) derived from the velocity field (v) and in most cases these vector fields differ fundamentally from each other. This phenomenon gains an immense significance for modelling some well-known fluids, e.g. animal blood or liquid crystals (see e.g. [PRU74]).

In the microscale, when at least one dimension of the domain is only a few times larger than the size of the molecules (e.g. blood vessels, lubricants), fluid motions even for isotropic fluid can differ substantially from what would follow from the computations based entirely on the Navier-Stokes equations (see [SASM02]). This behavior is caused by the dominance of the surface stresses over body forces. Although not all the aspects of physical experiments have been fully explained but it is justified to assume that the surface stresses and the internal degrees of freedom of particles are the deciding factors for properties of fluid motion.

It is worth mentioning, that apart from A. Eringen, other mathematicians and physicists have proposed numerous generalizations of the Navier-Stokes equations. The comparison of these theories can be found in [ATS73]. For a short historical review we refer the reader to [Łuk99, Ch. 1, §5].

In this work we plan to investigate the global existence of strong solutions to micropolar equations which are given by

$$
\begin{aligned}
&v_{,t} + v\cdot\nabla v - (\nu+\nu_r)\triangle v + \nabla p = 2\nu_r \operatorname{rot}\omega + f && \text{in } \Omega^\infty := \Omega\times(0,\infty),\\
&\operatorname{div} v = 0 && \text{in } \Omega^\infty,\\
&\omega_{,t} + v\cdot\nabla\omega - \alpha\triangle\omega - \beta\nabla\operatorname{div}\omega + 4\nu_r\omega = 2\nu_r\operatorname{rot} v + g && \text{in } \Omega^\infty,\\
&v|_{t=0} = v(0), \qquad \omega|_{t=0} = \omega(0) && \text{in } \Omega.
\end{aligned}
\tag{1.1}
$$

The unknowns are: the velocity field v, the pressure p and the microroation field ω. The viscosity coefficients ν, ν_r, α and β are fixed and positive. Note that if ν_r then $(1.1)_{1,2}$ and $(1.1)_3$ get uncoupled. Therefore we cannot expect better results than for the classical Navier-Stokes equations.

So far we have not specified the domain Ω. We assume that is has a product form

$$\left\{(x_1,x_2)\in\mathbb{R}^2\colon \varphi(x_1,x_2)\le c_0\right\}\times\left\{x_3\colon -a\le x_3\le a\right\},$$

where the constants a and c_0 are positive and φ is a $\mathcal{C}^2$-closed curve in $\mathbb{R}^2$. Thus, Ω is a finite cylinder placed alongside the x_3-axis (see Figure 1).

2000 Mathematics Subject Classification. 35Q30, 76D05.

Key words and phrases. micropolar fluids, cylindrical domains, global existence, strong solutions.

The author is partially supported by Polish KBN grant N N201 393137.



FIGURE 1. The domain Ω and its boundary $S := S_1 \cup S_2$.

From practical point of view (blood vessels, lubrication theory) our choice is justified. From theoretical perspective, our approach is intensely focused upon search for such solutions that are close to two dimensional (see e.g. [Zaj05], [RZ08], [Zaj11]). The solutions which are proved to exist, can be regarded as a slight perturbation of two dimensional flow along the perpendicular direction. This perturbation will be somehow measured by $\delta(t)$ (see (2.2)), which we introduce later.

We shall emphasize that since we only require the initial rate of change of the flow and microrotation, as well as the derivatives of the external data with respect to x_3 to be small, the flow alongside the cylinder can be large, but close to constant.

As far as the boundary condition are concerned, we use

$$
\begin{aligned}
&v \cdot n = 0 && \text{on } S^\infty := S \times (0,\infty),\\
&\operatorname{rot} v \times n = 0 && \text{on } S^\infty,\\
&\omega = 0 && \text{on } S_1^\infty,\\
&\omega' = 0, \qquad \omega_{3,x_3} = 0 && \text{on } S_2^\infty,
\end{aligned}
\tag{1.2}
$$

where n is the unit outward vector. The first two equations may be interpreted as tangential "slip" velocity being proportional to tangential stress with a factor of proportionality depending only on the curvature of φ (see e.g. [CMR98], [Kel06]). Such boundary condition was already postulated in 1827 by C.M.L.H. Navier. The third equation is clear but from the physical point view not necessarily adequate because the molecules may not move but they can rotate (see [BS70]). This effect is regarded in the fourth equation (see [Mig84]).

2. NOTATION

Before we present the main result of this work, we should employ some notation. By Ω^t we denote $\Omega \times (t_0, t)$ where $0 \le t_0 < t < \infty$. A generic constant c may change from line to line and is subscripted with appropriate symbol which indicates the dependence on the domain (c_Ω), embedding theorems (c_I), the Poincaré inequality (c_P) and viscosity coefficients ($c_{\alpha,\beta,\nu,\nu_r}$). To simplify the formulas we will also use

$$
h := v_{,x_3}, \qquad\qquad \theta := \omega_{,x_3}.
$$

To shorten energy estimate we introduce

$$
\begin{aligned}
E_{v,\omega}(t) &:= \|f\|_{L_2(t_0,t;L_{\frac{6}{5}}(\Omega))} + \|g\|_{L_2(t_0,t;L_{\frac{6}{5}}(\Omega))} + \|v(t_0)\|_{L_2(\Omega)} + \|\omega(t_0)\|_{L_2(\Omega)},\\
E_{h,\theta}(t) &:= \|f_{,x_3}\|_{L_2(t_0,t;L_{\frac{6}{5}}(\Omega))} + \|g_{,x_3}\|_{L_2(t_0,t;L_{\frac{6}{5}}(\Omega))} + \|h(t_0)\|_{L_2(\Omega)} + \|\theta(t_0)\|_{L_2(\Omega)}.
\end{aligned}
\tag{2.1}
$$

Finally, the function

$$
\delta(t) := \|f_{,x_3}\|^2_{L_2(\Omega^t)} + \|g_{,x_3}\|^2_{L_2(\Omega^t)} + \|\operatorname{rot} h(t_0)\|^2_{L_2(\Omega)} + \|h(t_0)\|^2_{L_2(\Omega)} + \|\theta(t_0)\|^2_{L_2(\Omega)}
\tag{2.2}
$$

will be of particular interest. It expresses the smallness assumption which has to be made in order to prove the existence of regular solutions on (t_0, t). Note that it contains only derivative of the external and the initial data with respect to x_3.

The notation for function spaces is standard and follows [Łuk99, Ch. 3, §1.1], [LSU67, Ch. 2, §3] and [Tem79, Ch. 1, §1.1]:

- $W^m_p(\Omega)$, where $m \in \mathbb{N}$, $p \geq 1$, is the closure of $\mathcal{C}^\infty(\Omega)$ in the norm

$$\|u\|_{W^m_p(\Omega)} = \left(\sum_{|\alpha| \leq m} \|\mathrm{D}^\alpha u\|^p_{L_p(\Omega)}\right)^{\frac{1}{p}},$$

- $H^k(\Omega)$, where $k \in \mathbb{N}$, is simply $W^k_2(\Omega)$,
- $W^{2,1}_p(\Omega^t)$, where $p \geq 1$, is the closure of $\mathcal{C}^\infty(\Omega \times (t_0, t_1))$ in the norm

$$\|u\|_{W^{2,1}_p(\Omega^t)} = \left(\int_{t_0}^{t_1}\int_\Omega |u_{,xx}(x,s)|^p + |u_{,x}(x,s)|^p + |u(x,s)|^p + |u_{,t}(x,s)|^p \,\mathrm{d}x\,\mathrm{d}s\right)^{\frac{1}{p}},$$

- $H^1_0(\Omega)$ is the closure of $\mathcal{C}^\infty_0(\Omega)$ in the norm

$$\|u\|_{H^1_0(\Omega)} = \left(\int_\Omega |\nabla u(x)|^2 \,\mathrm{d}x\right)^{\frac{1}{2}},$$

- $L_q(t_0, t_1; X)$, where $q \geq 1$ and X is a Banach space, is the set of all strongly measurable functions defined on the interval $[t_0, t_1]$ with values in X with finite norm defined by

$$\|u\|_{L_q(t_0,t_1;X)} = \left(\int_{t_0}^{t_1} \|u(s)\|^q_X \,\mathrm{d}s\right)^{\frac{1}{q}},$$

  where $1 \leq p < \infty$ and by

$$\|u\|_{L_\infty(t_0,t_1;X)} = \operatorname*{ess\,sup}_{t_0 \leq s \leq t_1} \|u(s)\|_X,$$

  for $q = \infty$.
- $V^k_2(\Omega^t)$, where $k \in \mathbb{N}$, is the closure of $\mathcal{C}^\infty(\Omega \times (t_0, t_1))$ in the norm

$$\|u\|_{V^k_2(\Omega^t)} = \operatorname*{ess\,sup}_{t \in (t_0,t_1)} \|u\|_{H^k(\Omega)} + \left(\int_{t_0}^{t_1} \|\nabla u\|^2_{H^k(\Omega)} \,\mathrm{d}t\right)^{1/2}$$

3. Main result

The main results of this work reads:

Theorem 1 (global existence)**.** *Let $t_0 = 0$ and $0 < T < \infty$ be sufficiently large and fixed. Suppose that $v(0), \omega(0) \in H^1(\Omega)$ and $\operatorname{rot} h(0) \in L_2(\Omega)$. In addition, let the external data satisfy $f_3|_{S_2} = 0$, $g'|_{S_2} = 0$,*

$$\|f(t)\|_{L_2(\Omega)} \leq \|f(kT)\|_{L_2(\Omega)} e^{-(t-kT)}, \qquad \|f_{,x_3}(t)\|_{L_2(\Omega)} \leq \|f_{,x_3}(kT)\|_{L_2(\Omega)} e^{-(t-kT)},$$
$$\|g(t)\|_{L_2(\Omega)} \leq \|g(kT)\|_{L_2(\Omega)} e^{-(t-kT)}, \qquad \|g_{,x_3}(t)\|_{L_2(\Omega)} \leq \|g_{,x_3}(kT)\|_{L_2(\Omega)} e^{-(t-kT)}$$

and

$$\sup_k \{f(kT), f_{,x_3}(kT), g(kT), g_{,x_3}(kT)\} < \infty.$$

Then, for $\delta(T)$ sufficiently small there exists a unique and regular solution to problem (1.1) *equipped with the boundary conditions* (1.2) *on the interval* $(0, \infty)$*. Moreover,*

$$\|v\|_{W^{2,1}_2(\Omega^\infty)} + \|\omega\|_{W^{2,1}_2(\Omega^\infty)} + \|\nabla p\|_{L_2(\Omega^\infty)} \leq \sup_k \Big(\|f\|_{L_2(\Omega^{kT})} + \|f_{,x_3}\|_{L_2(\Omega^{kT})}$$
$$+ \|g\|_{L_2(\Omega^{kT})} + \|g_{,x_3}\|_{L_2(\Omega^{kT})} + \|v(0)\|_{H^1(\Omega)} + \|\omega(0)\|_{H^1(\Omega)} + 1\Big)^3.$$

The proof of this theorem is based on the result obtained in [Now12, Theorem 1]:

Theorem 2 (large time existence)**.** *Let* $E_{v,\omega}(t) < \infty$*,* $E_{h,\theta}(t) < \infty$*. Suppose that* $v(t_0), \omega(t_0) \in H^1(\Omega)$*,* $f, g \in L_2(\Omega^t)$*. Finally, assume that* $f_3|_{S_2} = 0$*,* $g'|_{S_2} = 0$*. Then, for* $\delta(t)$ *sufficiently small there exists a unique solution* $(v,\omega) \in W_2^{2,1}(\Omega^t) \times W_2^{2,1}(\Omega^t)$ *to problem* (1.1) *supplemented with the boundary conditions* (1.2) *such that*

$$\|v\|_{W_2^{2,1}(\Omega^t)} + \|\nabla p\|_{L_2(\Omega^t)} \leq c_{\alpha,\nu,\nu_r,I,P,\Omega}\Big(E_{v,\omega}(t) + E_{h,\theta}(t) + \|f'\|_{L_2(\Omega^t)} + \|v(t_0)\|_{H^1(\Omega)} + 1\Big)^3$$

and

$$\begin{aligned}\|\omega\|_{W_2^{2,1}(\Omega^t)} \leq c_{\alpha,\nu,\nu_r,I,P,\Omega}\Big(&E_{v,\omega}(t) + E_{h,\theta}(t) + \|f'\|_{L_2(\Omega^t)} + \|g\|_{L_2(\Omega^t)} \\ &+ \|v(t_0)\|_{H^1(\Omega)} + \|\omega(t_0)\|_{H^1(\Omega)} + 1\Big)^3.\end{aligned}$$

We see that in view of Theorem 2 the extension of solutions up to the infinity with respect to time can be done in many ways. A substantive argument would be: if the solution to (1.1) on $[0,T]$ is regular and its estimate does not contain any constant which depend on time, then it is global. We must emphasize that the existing terminology on the topic is not precise. According to some authors a solution is global if the constants are time-dependent, but they do not blow up for any finite T. However, in such case it is more accurate to speak about large time existence instead of global existence.

For (1.1) we could not simply put $T = \infty$, because it would lead to improper integrals and several technical difficulties. Besides, it would imply that the external data must vanish as T goes to infinity. Hence, we adopt an alternative approach. We consider local solution on the time interval of the form $[kT,(k+1)T]$, where $k \in \mathbb{N}$ and $T > 0$ is fixed number. Starting with $k = 0$ we let $k \to \infty$, thereby obtaining a sequence of solutions with different initial conditions $v(kT)$ and $\omega(kT)$. In order to guarantee that this sequence is in fact an extension of solution from $[0,T]$, we must control the growth of the initial conditions. Additionally, if $f \neq 0$ and $g \neq 0$, certain restrictions on the external data must be also imposed.

Let us now briefly discuss previous results concerning the existence of global solutions of strong solutions to micropolar equations. In [Lan76] and [Lan77] Lange proved the existence and uniqueness of global and strong solutions to (1.1) in Hilbert spaces when the data are small enough. His approach is based on integral equations. This problem was also studied by Sava in [Sav78] under homogeneous boundary conditions in the case when the body forces and moments are not present. When the external data is present, ν is large and the data are small in comparison to ν problem (1.1) with zero Dirichlet boundary condition was examined by Łukaszewicz in [Łuk89] and in slightly more general framework by Ortega-Torres and Rojas-Medar in [OTRM97], who in contrast to Łukaszewicz did not assume any decay for the external data as t goes to the infinity. Semi-group approach was explored by Yamaguchi in [Yam05]. He established the global existence of strong solutions in case of small data.

For further bibliographical notes we refer the reader to [RM97], [OTRM97] and [Łuk99, Ch.3, §5].

Summing up: our result is established under smallness assumption not on the data itself but on its rate of change along x_3-variable. The external data do not vanish as t tends to infinity but exponentially decay on time intervals of the form $[kT,(k+1)T]$. The boundary conditions for the velocity field and partially for the microrotation field belong to the class of slip boundary conditions.

The rest of this work consists of Section 4, which contains some technical remarks, and Section 5 where we give estimates needed to prove Theorem 1.

4. Auxiliary tools

In this Section we present some technical facts which are essential for further considerations.

Lemma 4.1 (On integration by parts)**.** *Let* u *and* w *belong to* $H^1(\Omega)$*. Then*

$$\begin{aligned}\int_\Omega \operatorname{rot} u \cdot w \,\mathrm{d}x &= \int_\Omega \operatorname{rot} w \cdot u \,\mathrm{d}x + \int_S u \times n \cdot w \,\mathrm{d}S \\ &= \int_\Omega \operatorname{rot} w \cdot u \,\mathrm{d}x - \int_S w \times n \cdot u \,\mathrm{d}S\end{aligned}$$

Proof. It is an easy exercise. ☐

Lemma 4.2. *Suppose that*

$$\begin{aligned} \operatorname{rot} u &= \alpha \quad \text{in } \Omega, \\ \operatorname{div} u &= \beta \quad \text{in } \Omega \end{aligned}$$

with either $u \cdot n|_S = 0$ *or* $u \times n|_S = 0$. *Then*

$$\|u\|_{H^{k+1}(\Omega)} \le c_\Omega \big(\|\alpha\|_{H^k(\Omega)} + \|\beta\|_{H^k(\Omega)}\big).$$

Proof. For the proof we refer the reader [Sol73], where general overdetermined elliptic systems are examined. In particular, the case of tangent components of u is considered. ☐

Lemma 4.3. *Let* v, θ *and* $f_{,x_3}$ *be given. Then the pair* (h, q) *is a solution to the problem*

$$\begin{aligned} & h_{,t} - (\nu + \nu_r)\triangle h + \nabla q = -v \cdot \nabla h - h \cdot \nabla v + 2\nu_r \operatorname{rot} \theta + f_{,x_3} && \text{in } \Omega^t, \\ & \operatorname{div} h = 0 && \text{in } \Omega^t, \\ & \operatorname{rot} h \times n = 0, \quad h \cdot n = 0 && \text{on } S_1^t, \\ & h' = 0, \qquad h_{3,x_3} = 0 && \text{on } S_2^t, \\ & h|_{t=t_0} = h(t_0) && \text{in } \Omega. \end{aligned} \tag{4.1}$$

Lemma 4.4. *Let* h, ω, g' *and* $g_{,x_3}$ *be given. Then the function* θ *is solution to the problem*

$$\begin{aligned} & \theta_{,t} - \alpha\triangle\theta - \beta\nabla \operatorname{div}\theta + 4\nu_r\theta = -h \cdot \nabla\omega - v \cdot \nabla\theta + 2\nu_r \operatorname{rot} h + g_{,x_3} && \text{in } \Omega^t, \\ & \theta = 0 && \text{on } S_1^t, \\ & \theta_3 = 0, \qquad \theta'_{,x_3} = -\frac{1}{\alpha} g' && \text{on } S_2^t, \\ & \theta|_{t=t_0} = \theta(t_0) && \text{in } \Omega. \end{aligned} \tag{4.2}$$

Proofs of Lemmas 4.3 and 4.4 are given in [Now12].

Remark 4.5. Let us notice that for the functions h and θ the Poincaré inequality holds. Indeed, since h' vanishes on S_2 we only need to check if the integral of h_3 over Ω equals zero. We have

$$\int_\Omega h_3 \, \mathrm{d}x = \int_\Omega v_{3,x_3} \, \mathrm{d}x = \int_{S_2(x_3=-a)} v_3 \, \mathrm{d}x' - \int_{S_2(x_3=a)} v_3 \, \mathrm{d}x' = 0.$$

For θ_3, which vanishes on S_2 it is also obvious. For θ' we simply calculate the mean value:

$$\int_\Omega \theta' \, \mathrm{d}x = \int_\Omega \omega'_{,x_3} \, \mathrm{d}x = \int_{S_2(x_3=-a)} \omega' \, \mathrm{d}x' - \int_{S_2(x_3=a)} \omega' \, \mathrm{d}x' = 0,$$

which follows from $(1.2)_4$.

Lemma 4.6. *Suppose that* h *is a solution to* (4.1). *Then*

$$\|h\|_{H^2(\Omega)} \le c_\Omega \|\triangle h\|_{L_2(\Omega)}.$$

Proof. Consider

$$\begin{aligned} \operatorname{rot}\operatorname{rot} h &= \alpha && \text{in } \Omega, \\ \operatorname{div} h &= 0 && \text{in } \Omega, \\ \operatorname{rot} h \times n &= 0 && \text{on } S_1, \\ \operatorname{rot} h \cdot n &= 0 && \text{on } S_2. \end{aligned}$$

Introduce a partition of unity $\sum_{k=1}^N \zeta_k(x_3) = 1$. If we denote $\bar{u} = \operatorname{rot} h\zeta_k$, then the above system becomes

$$\begin{aligned} \operatorname{rot} \bar{u} &= \bar{\alpha} + 2\nabla h \cdot \nabla\zeta_k + h\triangle\zeta_k && \text{in } \operatorname{supp}\zeta_k \cap \Omega, \\ \operatorname{div} \bar{u} &= 0 && \text{in } \operatorname{supp}\zeta_k \cap \Omega, \\ \bar{u} \times n &= [0, 0, h_1\zeta_{k,x_3}n_1 + h_2\zeta_{k,x_3}n_2] && \text{on } \operatorname{supp}\zeta_k \cap S_1, \\ \bar{u} \cdot n &= 0 && \text{on } \operatorname{supp}\zeta_k \cap S_2. \end{aligned}$$

Note, that the boundary condition on $\operatorname{supp}\zeta_k \cap S_1$ is equal to zero which follows from $(4.1)_3$.

There are four cases to consider:

1. **$\operatorname{supp}\zeta_k \cap S = \emptyset$.:** The boundary conditions is $\bar u = 0$. From Lemma 4.2 we get

$$\|\bar u\|_{H^1(\operatorname{supp}\zeta_k\cap\Omega)} \le c_\Omega \left(\|\bar\alpha\|_{L_2(\operatorname{supp}\zeta_k\cap\Omega)} + \|\nabla h\|_{L_2(\operatorname{supp}\zeta_k\cap\Omega)} + \|h\|_{L_2(\operatorname{supp}\zeta_k\cap\Omega)}\right). \tag{4.3}$$

2. **$\operatorname{supp}\zeta_k \cap S_1 \neq \emptyset$, $\operatorname{supp}\zeta_k \cap S_2 = \emptyset$.:** On the set $\operatorname{supp}\zeta_k \cap S_1$ we see that $\bar u \times n = 0$, whereas on $\partial(\operatorname{supp}\zeta_k\cap\Omega) \setminus (\operatorname{supp}\zeta_k \cap S_1)$ we have $\bar u = 0$ which in particular means that $\bar u \times n = 0$. Next we transform the set $\operatorname{supp}\zeta_k\cap\Omega$ into the half-space and utilize Theorem 5.5 from [Sol65]. It yields (4.3) but in the half-space, i.e.

$$\|\bar u\|_{H^1(\mathbb{R}^{n-1}_+)} \le c_\Omega \left(\|\bar\alpha\|_{L_2(\mathbb{R}^{n-1}_+)} + \|\nabla h\|_{L_2(\mathbb{R}^{n-1}_+)} + \|h\|_{L_2(\mathbb{R}^{n-1}_+)}\right).$$

3. **$\operatorname{supp}\zeta_k \cap S_1 \neq \emptyset$, $\operatorname{supp}\zeta_k \cap S_2 \neq \emptyset$.:** Observe that $\bar u \cdot n|_{S_2} = \operatorname{rot}\bar h \cdot n|_{S_2} = 0 \Leftrightarrow h_{2,x_1} - h_{1,x_2} = 0$. On the other hand we already know that $h'|_{S_2} = 0$ and $h_{3,x_3} = 0$ on S_2 (see $(4.1)_4$). Therefore we can reflect the function h outside the cylinder following the formula

$$\check h(x) = \begin{cases} \bar h(x) & x_3 \in \overline{\operatorname{supp}\zeta_k\cap\Omega}, \\ (\bar h'(\bar x), -\bar h_3(\bar x)) & x_3 \le -a, \\ (\bar h'(\tilde x), -\bar h_3(\tilde x)) & x_3 \ge a, \end{cases}$$

   where $\bar x = (x', -2a - x_3)$ and $\tilde x = (x', 2a - x_3)$. We easily check that $\check u \times n = 0$ on $\operatorname{supp}\zeta_k \cap \check S_1$ and $\check u = 0$ on $\partial(\operatorname{supp}\zeta_k \cap \check\Omega)$. Then we follow Case 2.

4. **$\operatorname{supp}\zeta_k \cap S_1 = \emptyset$, $\operatorname{supp}\zeta_k \cap S_2 \neq \emptyset$.:** On $\operatorname{supp}\zeta_k\cap S_2$ we have $\bar u\cdot n = 0$ and on $\partial(\operatorname{supp}\zeta_k\cap\Omega)$ we get $\bar u = 0$. Next we map $\operatorname{supp}\zeta_k\cap\Omega$ into the half-space and utilize Theorem 5.5 from [Sol65]. It yields (4.3) in the half-space.

Summing over k yields

$$\|\operatorname{rot} h\|_{H^1(\Omega)} \le c_\Omega \left(\|\alpha\|_{L_2(\Omega)} + \|h\|_{H^1(\Omega)}\right) = c_\Omega \left(\|\triangle h\|_{L_2(\Omega)} + \|h\|_{H^1(\Omega)}\right).$$

From the above inequality and using (see [Now12, Rem. 8.3])

$$\|h\|_{H^{k+1}(\Omega)} \le c_\Omega \|\operatorname{rot} h\|_{H^k(\Omega)} \tag{4.4}$$

we deduce that for $\alpha = \operatorname{rot} h \in H^1(\Omega)$

$$\|h\|_{H^2(\Omega)} \le c_\Omega \|\alpha\|_{H^1(\Omega)} = c_\Omega \|\operatorname{rot} h\|_{H^1(\Omega)} \le c_\Omega \left(\|\triangle h\|_{L_2(\Omega)} + \|h\|_{H^1(\Omega)}\right).$$

Eventually, we demonstrate that $\|h\|_{H^1(\Omega)} \le \|\triangle h\|_{L_2(\Omega)}$. Again, from (4.4) it follows that

$$\|h\|^2_{H^1(\Omega)} \le c_\Omega \|\operatorname{rot} h\|^2_{L_2(\Omega)} = c_\Omega \int_\Omega \operatorname{rot} h \cdot \operatorname{rot} h \,\mathrm{d}x = \int_\Omega \operatorname{rot}\operatorname{rot} h \cdot h \,\mathrm{d}x - \int_S \operatorname{rot} h \times n \cdot h \,\mathrm{d}S,$$

where we also used Lemma 4.1. The boundary integral vanishes on S_1 due to boundary conditions $(4.1)_3$. On S_2 it can be written in a form

$$-\int_{S_2} h \times n \cdot \operatorname{rot} h \,\mathrm{d}x$$

and since $h \times n|_{S_2} = [-h_2, h_1, 0]|_{S_2} = 0$, where the last equality follows from $(4.1)_4$, we get

$$\|h\|^2_{H^1(\Omega)} \le c_\Omega \int_\Omega \operatorname{rot}\operatorname{rot} h \cdot h \,\mathrm{d}x = -c_\Omega \int_\Omega \triangle h \cdot h \,\mathrm{d}x.$$

Now we use the Hölder and the Young with ϵ inequalities, which results in

$$\|h\|^2_{H^1(\Omega)} \le c_\Omega \|\triangle h\|^2_{L_2(\Omega)}$$

and ends the proof. □

Lemma 4.7. *Let $E_{v,\omega}(t) < \infty$ (see $(2.1)_1$). Then for any $t_0 \le t \le t_1$ we have*

$$\|v\|_{V^0_2(\Omega^t)} + \|\omega\|_{V^0_2(\Omega^t)} \le c_{\alpha,\nu,I,\Omega} E_{v,\omega}(t).$$

Proof. This was proved in [Now12, Lemma 8.1]. □

5. Uniform estimates of solutions

We begin with certain refinement of the fundamental energy estimate for the function v and ω in the norm $L_\infty(t_0,t_1;L_2(\Omega))$.

Lemma 5.1. *Suppose that* $v(t_0),\omega(t_0)\in L_2(\Omega)$ *and* $f,g\in L_\infty(t_0,t;L_2(\Omega))$. *Then*

$$\begin{aligned}\|v(t)\|^2_{L_2(\Omega)}+\|\omega(t)\|^2_{L_2(\Omega)}&\le c_{\alpha,\nu,\Omega}\left(\|f\|^2_{L_\infty(t_0,t;L_2(\Omega))}+\|g\|^2_{L_\infty(t_0,t;L_2(\Omega))}\right)\\&\quad+\left(\|v(t_0)\|^2_{L_2(\Omega)}+\|\omega(t_0)\|^2_{L_2(\Omega)}\right)e^{-\frac{\min\{\alpha,\nu\}}{c_\Omega}(t-t_0)}.\end{aligned}\tag{A}$$

If, in addition we assume that

$$\begin{aligned}\|f(t)\|_{L_2(\Omega)}&\le\|f(t_0)\|_{L_2(\Omega)}e^{-(t-t_0)},\\\|g(t)\|_{L_2(\Omega)}&\le\|g(t_0)\|_{L_2(\Omega)}e^{-(t-t_0)}\end{aligned}$$

then (**A**) *implies*

$$\begin{aligned}\|v(t)\|^2_{L_2(\Omega)}+\|\omega(t)\|^2_{L_2(\Omega)}&\le\frac{c_\Omega}{\nu}\|f(t_0)\|^2_{L_2(\Omega)}e^{-(t-t_0)}+\frac{c_\Omega}{\alpha}\|g(t_0)\|^2_{L_2(\Omega)}e^{-(t-t_0)}\\&\quad+\left(\|v(t_0)\|^2_{L_2(\Omega)}+\|\omega(t_0)\|^2_{L_2(\Omega)}\right)e^{-\frac{\min\{\alpha,\nu\}}{c_\Omega}(t-t_0)}.\end{aligned}\tag{B}$$

To prove the above lemma we essentially follow the standard way of obtaining the basic energy estimates, however the final estimate is calculated in a slightly different way.

Proof. Multiplying $(1.1)_{1,2}$ by v and ω respectively and integrating over Ω yields

$$\begin{aligned}\frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}&\left(\|v\|^2_{L_2(\Omega)}+\|\omega\|^2_{L_2(\Omega)}\right)-(\nu+\nu_r)\int_\Omega\triangle v\cdot v\,\mathrm{d}x-\alpha\int_\Omega\triangle\omega\cdot\omega\,\mathrm{d}x-\beta\int_\Omega\nabla\operatorname{div}\omega\cdot\omega\,\mathrm{d}x\\&+\int_\Omega\nabla p\cdot v\,\mathrm{d}x+\int_\Omega v\cdot\nabla v\cdot v\,\mathrm{d}x+\int_\Omega v\cdot\nabla\omega\cdot\omega\,\mathrm{d}x+4\nu_r\|\omega\|^2_{L_2(\Omega)}\\&=2\nu_r\int_\Omega\operatorname{rot}\omega\cdot v\,\mathrm{d}x+2\nu_r\int_\Omega\operatorname{rot}v\cdot\omega\,\mathrm{d}x+\int_\Omega f\cdot v\,\mathrm{d}x+\int_\Omega g\cdot\omega\,\mathrm{d}x.\end{aligned}$$

In view of $(1.2)_1$ and from $\operatorname{div}v=0$ we immediately get that

$$\begin{aligned}\int_\Omega v\cdot\nabla v\cdot v\,\mathrm{d}x&=0,\\\int_\Omega v\cdot\nabla\omega\cdot\omega\,\mathrm{d}x&=0,\\\int_\Omega\nabla p\cdot v\,\mathrm{d}x&=0.\end{aligned}$$

By the application of Lemma 4.1 we see that

$$-\int_\Omega\triangle v\cdot v\,\mathrm{d}x=\int_\Omega\operatorname{rot}\operatorname{rot}v\cdot v\,\mathrm{d}x=\int_\Omega|\operatorname{rot}v|^2+\int_S\operatorname{rot}v\times n\cdot v\,\mathrm{d}S$$

and

$$\begin{aligned}-\int_\Omega\triangle\omega\cdot\omega,\mathrm{d}x=\int_\Omega\operatorname{rot}\operatorname{rot}\omega\cdot\omega-\nabla\operatorname{div}\omega\cdot\omega\,\mathrm{d}x&=\int_\Omega|\operatorname{rot}\omega|^2+|\operatorname{div}\omega|^2\,\mathrm{d}x\\&+\int_S\operatorname{rot}\omega\times n\cdot\omega\,\mathrm{d}S+\int_S\operatorname{div}\omega(\omega\cdot n)\,\mathrm{d}S.\end{aligned}$$

In view of the boundary conditions (1.2) all above boundary integrals vanish. On the right-hand side we have

$$\int_\Omega\operatorname{rot}\omega\cdot v\,\mathrm{d}x=\int_\Omega\operatorname{rot}v\cdot\omega\,\mathrm{d}x+\int_S\omega\times n\cdot v\,\mathrm{d}S,$$

where we applied Lemma 4.1. Next we see that the boundary integral vanishes on S_1 since $\omega|_{S_1} = 0$ and on S_2 $\omega \times n = [-\omega_2, \omega_1, 0] = 0$ holds, which follows from $(1.2)_4$. Thus

$$\begin{aligned}\frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}\left(\|v\|^2_{L_2(\Omega)} + \|\omega\|^2_{L_2(\Omega)}\right) + (\nu+\nu_r)\|\operatorname{rot} v\|^2_{L_2(\Omega)} + \alpha\|\operatorname{rot}\omega\|^2_{L_2(\Omega)} + (\alpha+\beta)\|\operatorname{div}\omega\|^2_{L_2(\Omega)} \\ + 4\nu_r\|\omega\|^2_{L_2(\Omega)} = 4\nu_r\int_\Omega \operatorname{rot} v\cdot\omega\,\mathrm{d}x + \int_\Omega f\cdot v\,\mathrm{d}x + \int_\Omega g\cdot\omega\,\mathrm{d}x.\end{aligned}$$

Now we make necessary estimates. From Lemma 4.2 for $u = \omega$ it follows that

$$\begin{aligned}\frac{\alpha}{c_\Omega}\|\omega\|^2_{H^1(\Omega)} &\le \alpha\left(\|\operatorname{rot}\omega\|^2_{L_2(\Omega)} + \|\operatorname{div}\omega\|^2_{L_2(\Omega)}\right),\\ \frac{\nu}{c_\Omega}\|v\|^2_{H^1(\Omega)} &\le \nu\|\operatorname{rot} v\|^2_{L_2(\Omega)}.\end{aligned}$$

By the Hölder and the Young inequalities we obtain

$$\begin{aligned}4\nu_r\int_\Omega \operatorname{rot} v\cdot\omega\,\mathrm{d}x &\le 4\nu_r\epsilon_1\|\operatorname{rot} v\|^2_{L_2(\Omega)} + \frac{\nu_r}{\epsilon_1}\|\omega\|^2_{L_2(\Omega)},\\ \int_\Omega f\cdot v\,\mathrm{d}x &\le \epsilon_2\|v\|^2_{L_2(\Omega)} + \frac{1}{4\epsilon_2}\|f\|^2_{L_2(\Omega)} \le \epsilon_2\|v\|^2_{H^1(\Omega)} + \frac{1}{4\epsilon_2}\|f\|^2_{L_2(\Omega)},\\ \int_\Omega g\cdot\omega\,\mathrm{d}x &\le \epsilon_3\|\omega\|^2_{L_2(\Omega)} + \frac{1}{4\epsilon_3}\|g\|^2_{L_2(\Omega)} \le \epsilon_3\|\omega\|^2_{H^1(\Omega)} + \frac{1}{4\epsilon_3}\|g\|^2_{L_2(\Omega)}.\end{aligned}$$

Now we set $\epsilon_1 = \frac{1}{4}$, $\epsilon_2 = \frac{\nu}{2c_\Omega}$, $\epsilon_3 = \frac{\alpha}{2c_\Omega}$. Since $\|v\|_{L_2(\Omega)} \le \|v\|_{H^1(\Omega)}$ and $\|\omega\|_{L_2(\Omega)} \le \|\omega\|_{H^1(\Omega)}$ we finally get

$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}\left(\|v\|^2_{L_2(\Omega)} + \|\omega\|^2_{L_2(\Omega)}\right) + \frac{\nu}{2c_\Omega}\|v\|^2_{L_2(\Omega)} + \frac{\alpha}{2c_\Omega}\|\omega\|^2_{L_2(\Omega)} \le \frac{c_\Omega}{2\nu}\|f\|^2_{L_2(\Omega)} + \frac{c_\Omega}{2\alpha}\|g\|^2_{L_2(\Omega)}.$$

With $\bar{c}_{\alpha,\nu,\Omega} = \frac{\min\{\alpha,\nu\}}{c_\Omega}$, the above inequality becomes

$$\frac{\mathrm{d}}{\mathrm{dt}}\left(\left(\|v\|^2_{L_2(\Omega)} + \|\omega\|^2_{L_2(\Omega)}\right)e^{\bar{c}_{\alpha,\nu,\Omega}t}\right) \le \frac{c_\Omega}{\nu}\|f\|^2_{L_2(\Omega)}e^{\bar{c}_{\alpha,\nu,\Omega}t} + \frac{c_\Omega}{\alpha}\|g\|^2_{L_2(\Omega)}e^{\bar{c}_{\alpha,\nu,\Omega}t}.$$

Integrating with respect to $t \in (t_0, t_1)$ yields

$$\begin{aligned}\left(\|v(t)\|^2_{L_2(\Omega)} + \|\omega(t)\|^2_{L_2(\Omega)}\right)e^{\bar{c}_{\alpha,\nu,\Omega}t} &\le \frac{c_\Omega}{\nu}\int_{t_0}^t \|f(s)\|^2_{L_2(\Omega)}e^{\bar{c}_{\alpha,\nu,\Omega}s}\,\mathrm{d}s\\ &+ \frac{c_\Omega}{\alpha}\int_{t_0}^t \|g(s)\|^2_{L_2(\Omega)}e^{\bar{c}_{\alpha,\nu,\Omega}s}\,\mathrm{d}s + \left(\|v(t_0)\|^2_{L_2(\Omega)} + \|\omega(t_0)\|^2_{L_2(\Omega)}\right)e^{\bar{c}_{\alpha,\nu,\Omega}t_0}\end{aligned}$$

or equivalently

$$\begin{aligned}\|v(t)\|^2_{L_2(\Omega)} + \|\omega(t)\|^2_{L_2(\Omega)} &\le \frac{c_\Omega}{\nu}e^{-\bar{c}_{\alpha,\nu,\Omega}t}\int_{t_0}^t \|f(s)\|^2_{L_2(\Omega)}e^{\bar{c}_{\alpha,\nu,\Omega}s}\,\mathrm{d}s\\ &+ \frac{c_\Omega}{\alpha}e^{-\bar{c}_{\alpha,\nu,\Omega}t}\int_{t_0}^t \|g(s)\|^2_{L_2(\Omega)}e^{\bar{c}_{\alpha,\nu,\Omega}s}\,\mathrm{d}s + \left(\|v(t_0)\|^2_{L_2(\Omega)} + \|\omega(t_0)\|^2_{L_2(\Omega)}\right)e^{-\bar{c}_{\alpha,\nu,\Omega}(t-t_0)},\end{aligned}$$

which proofs assertion (**A**).

Next we use the assumption on the external data in the above inequality

$$\begin{aligned}\frac{c_\Omega}{\nu}e^{-\bar{c}_{\alpha,\nu,\Omega}t}&\int_{t_0}^t \|f(s)\|^2_{L_2(\Omega)}e^{\bar{c}_{\alpha,\nu,\Omega}s}\,\mathrm{d}s + \frac{c_\Omega}{\alpha}e^{-\bar{c}_{\alpha,\nu,\Omega}t}\int_{t_0}^t \|g(s)\|^2_{L_2(\Omega)}e^{\bar{c}_{\alpha,\nu,\Omega}s}\,\mathrm{d}s\\ &\le \frac{c_\Omega}{\nu}e^{-\bar{c}_{\alpha,\nu,\Omega}t}\|f(t_0)\|^2_{L_2(\Omega)}\int_{t_0}^t e^{-(s-t_0)}e^{\bar{c}_{\alpha,\nu,\Omega}s}\,\mathrm{d}s\\ &+ \frac{c_\Omega}{\alpha}e^{-\bar{c}_{\alpha,\nu,\Omega}t}\|g(t_0)\|^2_{L_2(\Omega)}\int_{t_0}^t e^{-(s-t_0)}e^{\bar{c}_{\alpha,\nu,\Omega}s}\,\mathrm{d}s\\ &\le \frac{c_\Omega}{\nu}\|f(t_0)\|^2_{L_2(\Omega)}e^{-(t-t_0)} + \frac{c_\Omega}{\alpha}\|g(t_0)\|^2_{L_2(\Omega)}e^{-(t-t_0)}.\end{aligned}$$

Thus

$$\begin{aligned}\|v(t)\|^2_{L_2(\Omega)}+\|\omega(t)\|^2_{L_2(\Omega)}\leq \frac{c_\Omega}{\nu}\|f(t_0)\|^2_{L_2(\Omega)}e^{-(t-t_0)}+\frac{c_\Omega}{\alpha}\|g(t_0)\|^2_{L_2(\Omega)}e^{-(t-t_0)}\\ +\left(\|v(t_0)\|^2_{L_2(\Omega)}+\|\omega(t_0)\|^2_{L_2(\Omega)}\right)e^{-\bar{c}_{\alpha,\nu,\Omega}(t-t_0)},\end{aligned}$$

which is precisely assertion (**B**) of lemma. □

Lemma 5.2. *Suppose that* $v(t_0),\omega(t_0)\in H^1(\Omega)$. *Assume that*

$$\begin{aligned}&\|f(t)\|_{L_2(\Omega)}\leq\|f(t_0)\|_{L_2(\Omega)}e^{-(t-t_0)}, && \|f_{,x_3}(t)\|_{L_2(\Omega)}\leq\|f_{,x_3}(t_0)\|_{L_2(\Omega)}e^{-(t-t_0)},\\ &\|g(t)\|_{L_2(\Omega)}\leq\|g(t_0)\|_{L_2(\Omega)}e^{-(t-t_0)}, && \|g_{,x_3}(t)\|_{L_2(\Omega)}\leq\|g_{,x_3}(t_0)\|_{L_2(\Omega)}e^{-(t-t_0)}\end{aligned}$$

for $t_0\leq t\leq t_1$. *Then*

$$\|v(t_1)\|_{H^1(\Omega)}+\|\omega(t_1)\|_{H^1(\Omega)}\leq c_{\nu,\alpha,\beta,I,P,\Omega}(t_1)\left(\|v(t_0)\|_{H^1(\Omega)}+\|\omega(t_0)\|_{H^1(\Omega)}\right),$$

where

$$\lim_{t_1\to\infty}c_{\nu,\alpha,\beta,I,P,\Omega}(t_1)=0.$$

Remark 5.3. We have assumed an exponential decay with respect to time on the external data and their derivative along the axis of the cylinder. The reason underlying this assumption follows from the necessity to control the amount of the energy supplied to the system. If we consider the global in time existence, the supplied energy must be balanced by the rate of its lost due to friction, otherwise the system blows-up. However, we emphasize that this exponential decay with respect to time for the external data has only a local character. If t_1 denotes the end of the time interval under consideration, then

$$\lim_{t\to t_1^-}\|f(t)\|_{L_2(\Omega)}\neq\lim_{t\to t_1^+}\|f(t)\|_{L_2(\Omega)}$$

and analogously for the other external data. The right-hand side limit can be even much larger than the left-hand side limit.

Note, that alternatively we may attempt to analyze carefully the direct correspondence between the energy input and loss which is likely to result in different assumption on the external data. However, it is beyond the scope of our study. For further reading we refer the reader to [LL87].

Note also, that if we had the Poincaré inequality for f and g with respect to x_3, we would be able to relax the assumption on the exponential decay of the external data and limit our considerations only to their rate of change.

Proof. First we multiply $(1.1)_{1,2}$ by $-\alpha\triangle v$ and $-\alpha\triangle\omega-\beta\nabla\operatorname{div}\omega$ respectively and integrate over Ω. It yields

$$\begin{aligned}(5.1a)\qquad -\alpha\int_\Omega v_{,t}\cdot\triangle v\,\mathrm{d}x+(\nu+\nu_r)\alpha\|\triangle v\|^2_{L_2(\Omega)}-\alpha\int_\Omega v\cdot\nabla v\cdot\triangle v\,\mathrm{d}x-\alpha\int_\Omega\nabla p\cdot\triangle v\,\mathrm{d}x\\ =-2\nu_r\alpha\int_\Omega\operatorname{rot}\omega\cdot\triangle v\,\mathrm{d}x-\alpha\int_\Omega f\cdot\triangle v\,\mathrm{d}x\end{aligned}$$

and

$$\begin{aligned}(5.1b)\qquad &-\int_\Omega\omega_{,t}\cdot(\alpha\triangle\omega+\beta\nabla\operatorname{div}\omega)\,\mathrm{d}x+\int_\Omega(\alpha\triangle\omega+\beta\nabla\operatorname{div}\omega)^2\,\mathrm{d}x\\ &-4\nu_r\int_\Omega\omega\cdot(\alpha\triangle\omega+\beta\nabla\operatorname{div}\omega)\,\mathrm{d}x=-2\nu_r\int_\Omega\operatorname{rot}v\cdot(\alpha\triangle\omega+\beta\nabla\operatorname{div}\omega)\,\mathrm{d}x\\ &-\int_\Omega g\cdot(\alpha\triangle\omega+\beta\nabla\operatorname{div}\omega)\,\mathrm{d}x+\int_\Omega v\cdot\nabla\omega\cdot(\alpha\triangle\omega+\beta\nabla\operatorname{div}\omega)\,\mathrm{d}x.\end{aligned}$$

By application of Lemma 4.1 we see that

$$\begin{aligned}&-\alpha\int_\Omega v_{,t}\cdot\triangle v\,\mathrm{d}x=\alpha\int_\Omega\operatorname{rot}v_{,t}\cdot\operatorname{rot}v\,\mathrm{d}x+\alpha\int_S\operatorname{rot}v\times n\cdot v_{,t}\,\mathrm{d}S=\frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}\alpha\int_\Omega|\operatorname{rot}v|^2\,\mathrm{d}x,\\ &-\alpha\int_\Omega\nabla p\cdot\triangle v\,\mathrm{d}x=\alpha\int_\Omega\operatorname{rot}\nabla p\cdot\operatorname{rot}v\,\mathrm{d}x+\alpha\int_S\operatorname{rot}v\times n\cdot\nabla p\,\mathrm{d}S=0,\end{aligned}$$

where we used the boundary conditions $(1.2)_2$. From the vector identity $-\triangle\omega = \operatorname{rot}\operatorname{rot}\omega - \nabla\operatorname{div}\omega$ and Lemma 4.1 it follows that

$$\begin{aligned}-\int_\Omega \omega_{,t}\cdot\triangle\omega\,\mathrm{d}x &= \int_\Omega \omega_{,t}\cdot\operatorname{rot}\operatorname{rot}\omega - \nabla\operatorname{div}\omega\,\mathrm{d}x = \frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}\int_\Omega |\operatorname{rot}\omega|^2\,\mathrm{d}x - \int_S \omega_{,t}\times n\cdot\operatorname{rot}\omega\,\mathrm{d}S\\ &+ \int_\Omega \omega_{,t}\cdot\nabla\operatorname{div}\omega\,\mathrm{d}x = \frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}\int_\Omega |\operatorname{rot}\omega|^2 + |\operatorname{div}\omega|^2\,\mathrm{d}x - \int_S \operatorname{div}\omega\,(\omega_{,t}\cdot n)\,\mathrm{d}S.\end{aligned}$$

The boundary conditions $(1.2)_4$ imply that $\omega'_{,x'}|_{S_2} = 0$. Thus, $\operatorname{div}\omega|_{S_2} = 0$. Since $\omega|_{S_1} = 0 \Rightarrow \omega_{,t}|_{S_1} = 0$ and $\omega_{,t}\times n = [-\omega_{2,t},\omega_{1,t},0] = 0$ we see that the boundary integrals vanish. Therefore

$$-\int_\Omega \omega_{,t}\cdot(\alpha\triangle\omega + \beta\nabla\operatorname{div}\omega)\,\mathrm{d}x = \frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}\int_\Omega \alpha|\operatorname{rot}\omega|^2 + (\alpha+\beta)|\operatorname{div}\omega|^2\,\mathrm{d}x.$$

In the same manner we get

$$-\int_\Omega \omega\cdot(\alpha\triangle\omega + \beta\nabla\operatorname{div}\omega)\,\mathrm{d}x = \int_\Omega \alpha|\operatorname{rot}\omega|^2 + (\alpha+\beta)|\operatorname{div}\omega|^2\,\mathrm{d}x.$$

Moving to the first term on the right-hand side we encounter a problem with the integral

$$\beta\int_\Omega \operatorname{rot}v\cdot\nabla\operatorname{div}\omega\,\mathrm{d}x.$$

We easily see that the application of Lemma 4.1 or integration by parts lead to a boundary integral, either

$$\beta\int_S v\times n\cdot\nabla\operatorname{div}\omega\,\mathrm{d}S \qquad\text{or}\qquad \beta\int_S \operatorname{div}\omega\operatorname{rot}v\cdot n\,\mathrm{d}S.$$

We are not able to compute these integrals, because the boundary conditions (1.2) yield insufficient information. The first integral contains second-order derivatives of ω, which makes it impossible to estimate by the data in any suitable norm. For the second we apply trace theorem and interpolation inequality, which we shall present later. For now we only write

$$\begin{aligned}-\int_\Omega \operatorname{rot}v\cdot(\alpha\triangle\omega + \beta\nabla\operatorname{div}\omega)\,\mathrm{d}x &= \alpha\int_\Omega \operatorname{rot}v\cdot\operatorname{rot}\operatorname{rot}\omega\,\mathrm{d}x - (\alpha+\beta)\int_\Omega \operatorname{rot}v\cdot\nabla\operatorname{div}\omega\,\mathrm{d}x\\ &= -\alpha\int_\Omega \triangle v\cdot\operatorname{rot}\omega\,\mathrm{d}x - \alpha\int_S \operatorname{rot}v\times n\cdot\operatorname{rot}\omega\,\mathrm{d}S - (\alpha+\beta)\int_S \operatorname{div}\omega\operatorname{rot}v\cdot n\,\mathrm{d}S\\ &= -\alpha\int_\Omega \triangle v\cdot\operatorname{rot}\omega\,\mathrm{d}x - (\alpha+\beta)\int_S \operatorname{div}\omega\operatorname{rot}v\cdot n\,\mathrm{d}S.\end{aligned}$$

Finally, we rewrite (5.1) in the following form

$$\begin{aligned}&\frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}\left(\int_\Omega \alpha|\operatorname{rot}v|^2 + \alpha|\operatorname{rot}\omega|^2 + (\alpha+\beta)|\operatorname{div}\omega|^2\,\mathrm{d}x\right) + (\nu+\nu_r)\alpha\|\triangle v\|^2_{L_2(\Omega)}\\ &\quad + \|\alpha\triangle\omega + \beta\nabla\operatorname{div}\omega\|^2_{L_2(\Omega)} + 4\nu_r\alpha\int_\Omega |\operatorname{rot}\omega|^2 + (\alpha+\beta)|\operatorname{div}\omega|^2\,\mathrm{d}x\\ &= -4\nu_r\alpha\int_\Omega \operatorname{rot}\omega\cdot\triangle v\,\mathrm{d}x - 2\nu_r(\alpha+\beta)\int_S \operatorname{div}\omega\operatorname{rot}v\cdot n\,\mathrm{d}S + \int_\Omega v\cdot\nabla v\cdot\triangle v\,\mathrm{d}x\\ &\quad + \int_\Omega v\cdot\nabla\omega\cdot(\alpha\triangle\omega + \beta\nabla\operatorname{div}\omega)\,\mathrm{d}x - \int_\Omega f\cdot\triangle v\,\mathrm{d}x - \int_\Omega g\cdot(\alpha\triangle\omega + \beta\nabla\operatorname{div}\omega)\,\mathrm{d}x\\ &=: \sum_{k=1}^{6} I_k.\end{aligned}\tag{5.2}$$

We shall estimate I_k by application of the Hölder and the Young inequalities. Two first one is obvious:

$$I_1 \le 4\nu_r\alpha\|\operatorname{rot}\omega\|_{L_2(\Omega)}\|\triangle v\|_{L_2(\Omega)} \le 4\nu_r\alpha\epsilon_1\|\triangle v\|^2_{L_2(\Omega)} + \frac{\nu_r\alpha}{\epsilon_1}\|\operatorname{rot}\omega\|^2_{L_2(\Omega)}.$$

For I_2 more work is required. From the trace theorem it follows that for $\epsilon > 0$

$$\begin{aligned} I_2 &\le 2\nu_r(\alpha+\beta)\|\operatorname{div}\omega\|_{L_2(S)}\|\operatorname{rot} v\cdot n\|_{L_2(S)} \le \nu_r(\alpha+\beta)\|\operatorname{div}\omega\|^2_{L_2(S)} + \nu_r(\alpha+\beta)\|\operatorname{rot} v\|^2_{L_2(S)} \\ &\le c_\Omega\nu_r(\alpha+\beta)\|\operatorname{div}\omega\|^2_{H^{\frac12-\epsilon}(\Omega)} + c_\Omega(\alpha+\beta)\|\operatorname{rot} v\|^2_{H^{\frac12-\epsilon}(\Omega)}. \end{aligned}$$

Now we use an inequality of Gagliardo-Nirenberg type (see [CD00, Ch. 1, Rem. 1.2.1])

$$\begin{aligned} \|\operatorname{div}\omega\|^2_{H^{\frac12-\epsilon}(\Omega)} &\le \|\omega\|^{2\theta}_{H^2(\Omega)}\|\omega\|^{2(1-\theta)}_{L_2(\Omega)}, \\ \|\operatorname{rot} v\|^2_{H^{\frac12-\epsilon}(\Omega)} &\le \|v\|^{2\theta}_{H^2(\Omega)}\|v\|^{2(1-\theta)}_{L_2(\Omega)}, \end{aligned}$$

where θ satisfies

$$\frac32 - \epsilon - \frac32 \le (1-\theta)\left(-\frac32\right) + \theta\left(2-\frac32\right) \quad\Leftrightarrow\quad \frac32 - \epsilon \le 2\theta,$$

which allows us to set $\theta = \frac34$. Thus, by the Young inequality with ϵ

$$\begin{aligned} \|\operatorname{div}\omega\|^2_{H^{\frac12-\epsilon}(\Omega)} &\le \|\omega\|^{\frac32}_{H^2(\Omega)}\|\omega\|^{\frac12}_{L_2(\Omega)} \le \epsilon_{21}\|\omega\|^2_{H^2(\Omega)} + \frac{1}{4\epsilon_{21}}\|\omega\|^2_{L_2(\Omega)} \\ \|\operatorname{rot} v\|^2_{H^{\frac12-\epsilon}(\Omega)} &\le \|v\|^{\frac32}_{H^2(\Omega)}\|v\|^{\frac12}_{L_2(\Omega)} \le \epsilon_{22}\|v\|^2_{H^2(\Omega)} + \frac{1}{4\epsilon_{22}}\|v\|^2_{L_2(\Omega)}. \end{aligned}$$

Finally

$$\begin{aligned} I_2 \le c_\Omega\nu_r(\alpha+\beta)\epsilon_{21}\|\omega\|^2_{H^2(\Omega)} &+ \frac{c_\Omega\nu_r(\alpha+\beta)}{4\epsilon_{21}}\|\omega\|^2_{L_2(\Omega)} \\ &+ c_\Omega\nu_r(\alpha+\beta)\epsilon_{22}\|v\|^2_{H^2(\Omega)} + \frac{c_\Omega\nu_r(\alpha+\beta)}{4\epsilon_{22}}\|v\|^2_{L_2(\Omega)}. \end{aligned}$$

For the nonlinear terms we have

$$I_3 \le \|\triangle v\|_{L_2(\Omega)}\|\nabla v\|_{L_2(\Omega)}\|v\|_{L_\infty(\Omega)} \le \epsilon_3\|\triangle v\|^2_{L_2(\Omega)} + \frac{1}{4\epsilon_3}\|\nabla v\|^2_{L_2(\Omega)}\|v\|^2_{L_\infty(\Omega)}$$

and

$$\begin{aligned} I_4 &\le \|\alpha\triangle\omega + \beta\nabla\operatorname{div}\omega\|_{L_2(\Omega)}\|\nabla\omega\|_{L_2(\Omega)}\|v\|_{L_\infty(\Omega)} \\ &\le \epsilon_4(\alpha+\beta)\|\omega\|^2_{H^2(\Omega)} + \frac{\alpha+\beta}{4\epsilon_4}\|\nabla\omega\|^2_{L_2(\Omega)}\|v\|^2_{L_\infty(\Omega)}. \end{aligned}$$

The two last terms are estimated in a standard way

$$\begin{aligned} I_5 &\le \epsilon_5\|\triangle v\|^2_{L_2(\Omega)} + \frac{1}{4\epsilon_5}\|f\|^2_{L_2(\Omega)}, \\ I_6 &\le \epsilon_6(\alpha+\beta)\|\omega\|^2_{H^2(\Omega)} + \frac{\alpha+\beta}{4\epsilon_6}\|g\|^2_{L_2(\Omega)}. \end{aligned}$$

Before we chose ϵ_i, $i = 1,\dots,6$, we justify the following inequality

$$\|v\|^2_{H^2(\Omega)} \le c_\Omega\|\triangle v\|^2_{L_2(\Omega)}. \tag{5.3}$$

If we put $u = \operatorname{rot} v$ (we see that $u\times n|_S = 0$) in Lemma 4.1 then we get

$$\|\operatorname{rot} v\|^2_{H^1(\Omega)} \le c_\Omega\|\operatorname{rot}\operatorname{rot} v\|^2_{L_2(\Omega)} = c_\Omega\|\triangle v\|^2_{L_2(\Omega)}.$$

On the other hand, let now $u = v$ (we see that $u\cdot n|_S = 0$) in Lemma 4.1 and use the above inequality. Then

$$\|v\|^2_{H^2(\Omega)} \le c_\Omega\|\operatorname{rot} v\|^2_{H^1(\Omega)} \le c_\Omega\|\triangle v\|^2_{L_2(\Omega)}.$$

We set

$$\begin{aligned} &\epsilon_1 = \frac{1}{4\alpha}, \\ &\epsilon_{21}c_\Omega\nu_r(\alpha+\beta) = \epsilon_4(\alpha+\beta) = \epsilon_6(\alpha+\beta) = \frac{1}{6(c_\Omega(\alpha+\beta)+1)}, \\ &\epsilon_{22}c_\Omega^2\nu_r(\alpha+\beta) = \epsilon_3 = \epsilon_5 = \frac{\nu\alpha}{6}. \end{aligned}$$

Thus, we get from (5.2) that

$$\begin{aligned}\frac{1}{2}\left(\frac{\mathrm{d}}{\mathrm{dt}}\int_\Omega \alpha|\operatorname{rot} v|^2 + \alpha|\operatorname{rot}\omega|^2 + (\alpha+\beta)|\operatorname{div}\omega|^2\,\mathrm{d}x\right) &+ \frac{\nu\alpha}{2c_\Omega}\|v\|^2_{H^2(\Omega)} \\ &+ \frac{1}{2(c_\Omega(\alpha+\beta)+1)}\|\omega\|^2_{H^2(\Omega)} \\ \leq \frac{3c_\Omega^2\nu_r^2(\alpha+\beta)^2(c_\Omega(\alpha+\beta)+1)}{2}\|\omega\|^2_{L_2(\Omega)} &+ \frac{3c_\Omega^3\nu_r^2(\alpha+\beta)^2}{2\nu\alpha}\|v\|^2_{L_2(\Omega)} \\ + \frac{3}{2\nu\alpha}\|\nabla v\|^2_{L_2(\Omega)}\|v\|^2_{L_\infty(\Omega)} &+ \frac{3(c_\Omega(\alpha+\beta)+1)(\alpha+\beta)^2}{2}\|\nabla\omega\|^2_{L_2(\Omega)}\|v\|^2_{L_\infty(\Omega)} \\ &+ \frac{3}{2\nu\alpha}\|f\|^2_{L_2(\Omega)} + \frac{3(c_\Omega(\alpha+\beta)+1)(\alpha+\beta)^2}{2}\|g\|^2_{L_2(\Omega)}.\end{aligned}\tag{5.4}$$

where we use (5.3) at the end. The next step is to estimate the $H^2(\Omega)$-norms on the left-hand side from below by $L_2(\Omega)$-norms of $\operatorname{rot} v$ and $\operatorname{rot}\omega + \operatorname{div}\omega$. For any $u \in H^2(\Omega)$ we have

$$\frac{1}{2}\left(\|\operatorname{rot} u\|^2_{L_2(\Omega)} + \|\operatorname{div} u\|^2_{L_2(\Omega)}\right) \leq \|u\|^2_{H^1(\Omega)} \leq \|u\|^2_{H^2(\Omega)}.$$

Let

$$c_1 = \min\left\{\frac{\nu\alpha}{c_\Omega}, \frac{1}{(c_\Omega(\alpha+\beta)+1)}\right\}.$$

Then

$$\begin{aligned}\frac{\nu\alpha}{2c_\Omega}\|v\|^2_{H^2(\Omega)} + \frac{1}{2(c_\Omega(\alpha+\beta)+1)}\|\omega\|^2_{H^2(\Omega)} &\geq \frac{c_1}{2}\left(\|v\|^2_{H^2(\Omega)} + \|\omega\|^2_{H^2(\Omega)}\right) \\ &\geq \frac{c_1}{4}\left(\|\operatorname{rot} v\|^2_{L_2(\Omega)} + \|\operatorname{rot}\omega\|^2_{L_2(\Omega)} + \|\operatorname{div}\omega\|^2_{L_2(\Omega)}\right) \\ &\geq \frac{c_1}{4(\alpha+\beta)+2\delta}\left(\alpha\|\operatorname{rot} v\|^2_{L_2(\Omega)} + \beta\|\operatorname{rot}\omega\|^2_{L_2(\Omega)} + (\alpha+\beta)\|\operatorname{div}\omega\|^2_{L_2(\Omega)}\right),\end{aligned}\tag{5.5}$$

where $\delta \geq 1$ will be chosen later. On the other hand, in view of Lemma 4.2 we already know that

$$\begin{aligned}\|\nabla v\|^2_{L_2(\Omega)} &\leq \|v\|^2_{H^1(\Omega)} \leq c_\Omega\|\operatorname{rot} v\|^2_{L_2(\Omega)}, \\ \|\nabla\omega\|^2_{L_2(\Omega)} &\leq \|\omega\|^2_{H^1(\Omega)} \leq c_\Omega\left(\|\operatorname{rot}\omega\|^2_{L_2(\Omega)} + \|\operatorname{div}\omega\|^2_{L_2(\Omega)}\right).\end{aligned}$$

Let

$$\begin{aligned}c_2 &= \max\left\{\frac{3}{\nu\alpha}, 3(c_\Omega(\alpha+\beta)+1)(\alpha+\beta)^2\right\}, \\ c_3 &= \max\left\{3c_\Omega^2\nu_r^2(\alpha+\beta)^2(c_\Omega(\alpha+\beta)+1), \frac{3c_\Omega^3\nu_r^2(\alpha+\beta)^2}{\nu\alpha}, \frac{3}{\nu\alpha}, 3(c_\Omega(\alpha+\beta)+1)(\alpha+\beta)^2\right\}.\end{aligned}$$

Denote

$$X(t) = \alpha\|\operatorname{rot} v(t)\|^2_{L_2(\Omega)} + \alpha\|\operatorname{rot}\omega(t)\|^2_{L_2(\Omega)} + (\alpha+\beta)\|\operatorname{div}\omega(t)\|^2_{L_2(\Omega)}.$$

Then, combining (5.5) with (5.4) gives

$$\begin{aligned}\frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}X(t) &+ \frac{c_1}{4(\alpha+\beta)+2\delta}X(t) \\ &\leq \frac{c_2}{2}\|v\|^2_{L_\infty(\Omega)}X(t) + \frac{c_3}{2}\left(\|f\|^2_{L_2(\Omega)} + \|g\|^2_{L_2(\Omega)} + \|v\|^2_{L_2(\Omega)} + \|\omega\|^2_{L_2(\Omega)}\right),\end{aligned}$$

which is equivalent to

$$\begin{aligned}\frac{\mathrm{d}}{\mathrm{dt}}X(t) &+ \left(\frac{c_1}{2(\alpha+\beta)+\delta} - c_2\|v\|^2_{L_\infty(\Omega)}\right)X(t) \\ &\leq c_3\left(\|f\|^2_{L_2(\Omega)} + \|g\|^2_{L_2(\Omega)} + \|v\|^2_{L_2(\Omega)} + \|\omega\|^2_{L_2(\Omega)}\right).\end{aligned}$$

We may also write

$$\begin{aligned}&\frac{\mathrm{d}}{\mathrm{dt}}\left(X(t)e^{\frac{c_1}{2(\alpha+\beta)+\delta}t-c_2\int_{t_0}^{t}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}\right)\\&\qquad\le c_3\left(\|f\|^2_{L_2(\Omega)}+\|g\|^2_{L_2(\Omega)}+\|v\|^2_{L_2(\Omega)}+\|\omega\|^2_{L_2(\Omega)}\right)e^{\frac{c_1}{2(\alpha+\beta)+\delta}t-c_2\int_{t_0}^{t}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}.\end{aligned}$$

Integrating with respect to $t\in(t_0,t_1)$ yields

$$\begin{aligned}&X(t_1)e^{\frac{c_1}{2(\alpha+\beta)+\delta}t_1-c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}\\&\qquad\le c_3\int_{t_0}^{t_1}\left(\|f(t)\|^2_{L_2(\Omega)}+\|g(t)\|^2_{L_2(\Omega)}+\|v(t)\|^2_{L_2(\Omega)}+\|\omega(t)\|^2_{L_2(\Omega)}\right)\\&\qquad\qquad\cdot e^{\frac{c_1}{2(\alpha+\beta)+\delta}t-c_2\int_{t_0}^{t}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}\,\mathrm{d}t+X(t_0)e^{\frac{c_1}{2(\alpha+\beta)+\delta}t_0}.\end{aligned}$$

Next we divide both sides by $e^{\frac{c_1}{2(\alpha+\beta)+\delta}t_1-c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}$. It gives

$$\begin{aligned}(5.6)\quad &X(t_1)\le c_3e^{-\frac{c_1}{2(\alpha+\beta)+\delta}t_1+c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}\\&\cdot\int_{t_0}^{t_1}\left(\|f(t)\|^2_{L_2(\Omega)}+\|g(t)\|^2_{L_2(\Omega)}+\|v(t)\|^2_{L_2(\Omega)}+\|\omega(t)\|^2_{L_2(\Omega)}\right)\cdot e^{\frac{c_1}{2(\alpha+\beta)+\delta}t-c_2\int_{t_0}^{t}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}\,\mathrm{d}t\\&\qquad+X(t_0)e^{-\frac{c_1}{2(\alpha+\beta)+\delta}(t_1-t_0)+c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}.\end{aligned}$$

Consider the integral with respect to t on the right-hand side. Since $\|v(s)\|^2_{L_\infty(\Omega)}$ is non-negative, we can write

$$e^{\frac{c_1}{2(\alpha+\beta)+\delta}t-c_2\int_{t_0}^{t}\|v(s)\|^2_{L_\infty(\Omega)}}\le e^{\frac{c_1}{2(\alpha+\beta)+\delta}t}.$$

By assumption and assertion (**B**) from Lemma 5.1 we see that

$$\begin{aligned}(5.7)\quad &c_3e^{-\frac{c_1}{2(\alpha+\beta)+\delta}t_1+c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}\\&\cdot\int_{t_0}^{t_1}\left(\|f(t)\|^2_{L_2(\Omega)}+\|g(t)\|^2_{L_2(\Omega)}+\|v(t)\|^2_{L_2(\Omega)}+\|\omega(t)\|^2_{L_2(\Omega)}\right)\cdot e^{\frac{c_1}{2(\alpha+\beta)+\delta}t-c_2\int_{t_0}^{t}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}\,\mathrm{d}t\\&\qquad\le c_3e^{-\frac{c_1}{2(\alpha+\beta)+\delta}t_1+c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}\\&\qquad\cdot\int_{t_0}^{t_1}\left(\|f(t)\|^2_{L_2(\Omega)}+\|g(t)\|^2_{L_2(\Omega)}+\|v(t)\|^2_{L_2(\Omega)}+\|\omega(t)\|^2_{L_2(\Omega)}\right)\cdot e^{\frac{c_1}{2(\alpha+\beta)+\delta}t}\,\mathrm{d}t\\&\qquad\le c_3e^{-\frac{c_1}{2(\alpha+\beta)+\delta}t_1+c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}\\&\qquad\cdot\left(\left(1+\frac{c_\Omega}{\nu}\right)\|f(t_0)\|^2_{L_2(\Omega)}+\left(1+\frac{c_\Omega}{\alpha}\right)\|g(t_0)\|^2_{L_2(\Omega)}+\|v(t_0)\|^2_{L_2(\Omega)}+\|\omega(t_0)\|^2_{L_2(\Omega)}\right)\\&\qquad\qquad\cdot\int_{t_0}^{t_1}e^{\frac{c_1}{2(\alpha+\beta)+\delta}t}\cdot e^{-\min\left\{1,\frac{\min\{\alpha,\nu\}}{c_\Omega}\right\}(t-t_0)}\,\mathrm{d}t.\end{aligned}$$

Now we chose δ. Two cases may occur:

1. $\alpha<c_\Omega\vee\nu<c_\Omega$.**:** In this case

$$\min\left\{1,\frac{\min\{\alpha,\nu\}}{c_\Omega}\right\}=\frac{\min\{\alpha,\nu\}}{c_\Omega},$$

so we chose such δ that it satisfies

$$\frac{c_1}{2(\alpha+\beta)+\delta}-\frac{\min\{\alpha,\nu\}}{c_\Omega}<0\quad\Leftrightarrow\quad\frac{c_1c_\Omega-2\min\{\alpha,\nu\}(\alpha+\beta)}{\min\{\alpha,\nu\}}<\delta.\tag{5.8}$$

Then

$$\begin{aligned}
&c_3 e^{-\frac{c_1}{2(\alpha+\beta)+\delta}t_1 + c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s} \int_{t_0}^{t_1} e^{\frac{c_1}{2(\alpha+\beta)+\delta}t}\cdot e^{-\min\left\{1,\frac{\min\{\alpha,\nu\}}{c_\Omega}\right\}(t-t_0)}\,\mathrm{d}t \\
&\quad = c_3 e^{-\frac{c_1}{2(\alpha+\beta)+\delta}t_1 + c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s + \frac{\min\{\alpha,\nu\}}{c_\Omega}t_0} \frac{1}{\frac{c_1}{2(\alpha+\beta)+\delta} - \frac{\min\{\alpha,\nu\}}{c_\Omega}} e^{\left(\frac{c_1}{2(\alpha+\beta)+\delta} - \frac{\min\{\alpha,\nu\}}{c_\Omega}\right)t}\Bigg|_{t_0}^{t_1} \\
&\quad \le c_3 e^{-\frac{c_1}{2(\alpha+\beta)+\delta}(t_1-t_0) + c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s} \frac{1}{\frac{\min\{\alpha,\nu\}}{c_\Omega} - \frac{c_1}{2(\alpha+\beta)+\delta}}.
\end{aligned}$$

2. **$\alpha > c_\Omega \wedge \nu > c_\Omega$.:** Now we have

$$\min\left\{1, \frac{\min\{\alpha,\nu\}}{c_\Omega}\right\} = 1.$$

Since $c_\Omega(\alpha+\beta)+1 > 1$ we see that $c_1 < 1$. It is enough to set $\delta = 1$. Then

$$\frac{c_1}{2(\alpha+\beta)+1} - 1 < 0$$

and

$$\begin{aligned}
&c_3 e^{-\frac{c_1}{2(\alpha+\beta)+\delta}t_1 + c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s} \int_{t_0}^{t_1} e^{\frac{c_1}{2(\alpha+\beta)+\delta}t}\cdot e^{-\min\left\{1,\frac{\min\{\alpha,\nu\}}{c_\Omega}\right\}(t-t_0)}\,\mathrm{d}t \\
&\quad = c_3 e^{-\frac{c_1}{2(\alpha+\beta)+1}t_1 + c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s + t_0} \frac{1}{\frac{c_1}{2(\alpha+\beta)+1} - 1} e^{\left(\frac{c_1}{2(\alpha+\beta)+1} - 1\right)t}\Bigg|_{t_0}^{t_1} \\
&\quad \le c_3 e^{-\frac{c_1}{2(\alpha+\beta)+1}(t_1-t_0) + c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s} \frac{1}{1 - \frac{c_1}{2(\alpha+\beta)+1}}.
\end{aligned}$$

In both cases we have obtained an estimate of the form

$$c_{\nu,\alpha,\beta,\Omega} c_3 e^{-\frac{c_1}{2(\alpha+\beta)+\delta}(t_1-t_0) + c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s},$$

where either $\delta = 1$ or δ satisfies (5.8). Putting the above estimate into (5.7), we get from (5.6) that

$$\begin{aligned}
X(t_1) &\le c_{\nu,\alpha,\beta,\Omega} c_3 e^{-\frac{c_1}{2(\alpha+\beta)+\delta}(t_1-t_0) + c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s} \\
&\quad \cdot c_{\nu,\alpha,\Omega}\left(\|f(t_0)\|^2_{L_2(\Omega)} + \|g(t_0)\|^2_{L_2(\Omega)} + \|v(t_0)\|^2_{L_2(\Omega)} + \|\omega(t_0)\|^2_{L_2(\Omega)}\right) \\
&\quad + X(t_0) e^{-\frac{c_1}{2(\alpha+\beta)+\delta}(t_1-t_0) + c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}.
\end{aligned}$$

In view of Lemma 4.2 and an obvious inequality $\alpha < \alpha+\beta$ we can rewrite the above estimate in the form

$$\begin{aligned}
\frac{\alpha}{c_\Omega}\left(\|v(t_1)\|^2_{H^1(\Omega)} + \|\omega(t_1)\|^2_{H^1(\Omega)}\right) &\le c_{\nu,\alpha,\beta,\Omega} c_3 e^{-\frac{c_1}{2(\alpha+\beta)+\delta}(t_1-t_0) + c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s} \\
&\quad \cdot c_{\nu,\alpha,\Omega}\left(\|f(t_0)\|^2_{L_2(\Omega)} + \|g(t_0)\|^2_{L_2(\Omega)} + \|v(t_0)\|^2_{L_2(\Omega)} + \|\omega(t_0)\|^2_{L_2(\Omega)}\right) \\
&\quad + 2(\alpha+\beta)\left(\|v(t_0)\|^2_{H^1(\Omega)} + \|\omega(t_0)\|^2_{H^1(\Omega)}\right) e^{-\frac{c_1}{2(\alpha+\beta)+\delta}(t_1-t_0) + c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s}.
\end{aligned}$$

Next observe that since $H^2(\Omega) \hookrightarrow L_\infty(\Omega)$, we have

$$\|v(s)\|^2_{L_\infty(\Omega)} \le c_I \|v(s)\|^2_{H^2(\Omega)}.$$

Integrating this inequality with respect to $t \in (t_0, t_1)$ and utilizing the estimate from Theorem 2 we obtain

$$\begin{aligned}
&\int_{t_0}^{t_1} \|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s \le c_I \|v\|^2_{L_2(t_0,t_1;H^2(\Omega))} \le c_I \|v\|^2_{W_2^{2,1}(\Omega^{t_1})} \\
&\le c_{\alpha,\nu,\nu_r,I,P,\Omega}\Big(E^3_{v,\omega}(t_1) + E^3_{h,\theta}(t_1) + \|f'\|^3_{L_2(\Omega^{t_1})} + \|v(t_0)\|^3_{H^1(\Omega)} + E_{v,\omega}(t_1) + \|v(t_0)\|_{H^1(\Omega)}\Big)^2.
\end{aligned}$$

By assumption on f, g, their derivative with respect to x_3 we see that

$$E_{v,\omega}(t_1)+E_{h,\theta}(t_1)\leq c_\Omega\Big(\|v(t_0)\|_{H^1(\Omega)}+\|\omega(t_0)\|_{H^1(\Omega)}+\|f(t_0)\|_{L_2(\Omega)}+\|g(t_0)\|_{L_2(\Omega)}\\ +\|f_{,x_3}(t_0)\|_{L_2(\Omega)}+\|g_{,x_3}(t_0)\|_{L_2(\Omega)}\Big).$$

Thus

$$\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s\leq\big(\|v(t_0)\|_{H^1(\Omega)}+\|\omega(t_0)\|_{H^1(\Omega)}+\|f(t_0)\|_{L_2(\Omega)}+\|g(t_0)\|_{L_2(\Omega)}\\ +\|f_{,x_3}(t_0)\|_{L_2(\Omega)}+\|g_{,x_3}(t_0)\|_{L_2(\Omega)}+1\big)^6$$

and therefore for t_1 large enough we have

$$-\frac{c_1}{2(\alpha+\beta)+\delta}(t_1-t_0)+c_2\int_{t_0}^{t_1}\|v(s)\|^2_{L_\infty(\Omega)}\,\mathrm{d}s<0,$$

which implies that

$$\|v(t_1)\|_{H^1(\Omega)}+\|\omega(t_1)\|_{H^1(\Omega)}\leq c_{\nu,\alpha,\beta,I,P,\Omega}(t_1)\left(\|v(t_0)\|_{H^1(\Omega)}+\|\omega(t_0)\|_{H^1(\Omega)}\right),$$

where the function $c_{\nu,\alpha,\beta,I,P,\Omega}(t_1)$ has the property that $\lim_{t_1\to\infty}c_{\nu,\alpha,\beta,I,P,\Omega}(t_1)=0$. This concludes the proof. □

Lemma 5.4. *Suppose that* $\operatorname{rot}h(t_0),h(t_0),\omega(t_0)\in H^1(\Omega)$, $f_3|_{S_2}=0$, $g'|_{S_2}=0$. *Assume that*

$$\|f(t)\|_{L_2(\Omega)}\leq\|f(t_0)\|_{L_2(\Omega)}e^{-(t-t_0)},\qquad \|f_{,x_3}(t)\|_{L_2(\Omega)}\leq\|f_{,x_3}(t_0)\|_{L_2(\Omega)}e^{-(t-t_0)},$$
$$\|g(t)\|_{L_2(\Omega)}\leq\|g(t_0)\|_{L_2(\Omega)}e^{-(t-t_0)},\qquad \|g_{,x_3}(t)\|_{L_2(\Omega)}\leq\|g_{,x_3}(t_0)\|_{L_2(\Omega)}e^{-(t-t_0)}$$

for any $t_0\leq t\leq t_1$. *Then*

$$\|\operatorname{rot}h(t_1)\|^2_{L_2(\Omega)}+\|h(t_1)\|^2_{L_2(\Omega)}+\|\theta(t_1)\|^2_{L_2(\Omega)}\\ \leq\|\operatorname{rot}h(t_0)\|^2_{L_2(\Omega)}+\|h(t_0)\|^2_{L_2(\Omega)}+\|\theta(t_0)\|^2_{L_2(\Omega)}.$$

Note, that we have assumed that the f_3 and g' vanish on S_2. Both functions appear in boundary integrals which need to be estimated. Although it is possible, but it leads to the presence of $c_I\left(\|h\|^2_{L_2(\Omega)}+\|\theta\|^2_{L_2(\Omega)}\right)$ on the right-hand side. Since c_I is out of control we are unable to ensure the uniform estimates analogously as we did in Lemma 5.1.

Proof of Lemma 5.4. We multiply $(4.1)_1$ by $-\triangle h$ and integrate over Ω, which yields

$$\begin{aligned}(5.9)\quad &-\int_\Omega h_{,t}\cdot\triangle h\,\mathrm{d}x+(\nu+\nu_r)\int_\Omega|\triangle h|^2\,\mathrm{d}x-\int_\Omega\nabla q\cdot\triangle h\,\mathrm{d}x\\ &=\int_\Omega v\cdot\nabla h\cdot\triangle h\,\mathrm{d}x+\int_\Omega h\cdot\nabla v\cdot\triangle h\,\mathrm{d}x-2\nu_r\int_\Omega\operatorname{rot}\theta\cdot\triangle h\,\mathrm{d}x-\int_\Omega f_{,x_3}\cdot\triangle h\,\mathrm{d}x.\end{aligned}$$

For the first term on the left-hand side we have

$$\begin{aligned}-\int_\Omega h_{,t}\cdot\triangle h\,\mathrm{d}x&=\int_\Omega h_{,t}\cdot\operatorname{rot}\operatorname{rot}h\,\mathrm{d}x\\ &=\frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}\int_\Omega|\operatorname{rot}h|^2\,\mathrm{d}x+\int_{S_1}h_{,t}\cdot\operatorname{rot}h\times n\,\mathrm{d}S_1+\int_{S_2}h_{2,t}\,(\operatorname{rot}h)_1-h_{1,t}\,(\operatorname{rot}h)_2\,\mathrm{d}S_2\\ &=\frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}\int_\Omega|\operatorname{rot}h|^2\,\mathrm{d}x\end{aligned}$$

where the boundary integrals vanish due to the boundary conditions $(4.1)_{3,4}$.

The third term on the left-hand side in (5.9) is equal to

$$\int_\Omega\nabla q\cdot\operatorname{rot}\operatorname{rot}h\,\mathrm{d}x=\int_\Omega\operatorname{rot}\nabla q\cdot\operatorname{rot}h\,\mathrm{d}x+\int_{S_1}\nabla q\cdot\operatorname{rot}h\times n\,\mathrm{d}S_1\\ +\int_{S_2}q_{,x_2}\,(\operatorname{rot}h)_1-q_{,x_1}\,(\operatorname{rot}h)_2\,\mathrm{d}S_2=0,$$

which follows from $(4.1)_3$ and

$$q|_{S_2} = -v\cdot\nabla v\cdot n|_{S_2} + f_3|_{S_2} + 2\nu_r \operatorname{rot}\omega\cdot n|_{S_2} = f_3|_{S_2} = 0$$

because we assumed that $f_3|_{S_2} = 0$.

Consider next the first term on the right-hand side in (5.9). Since $\operatorname{div} v = 0$ we may integrate by parts, which yields

$$\begin{aligned}\int_\Omega v\cdot\nabla h\cdot\triangle h\,\mathrm{d}x &\leq \|\nabla v\|_{L_6(\Omega)}\|\nabla h\|_{L_3(\Omega)}\|\operatorname{rot} h\|_{L_2(\Omega)} + \int_S (v\cdot\nabla)\,h\cdot(\operatorname{rot} h\times n)\,\mathrm{d}S\\ &\leq \epsilon_1 c_I\|\nabla h\|^2_{H^1(\Omega)} + \frac{1}{4\epsilon_1}\|\nabla v\|^2_{L_6(\Omega)}\|\operatorname{rot} h\|^2_{L_2(\Omega)} \leq \epsilon_1 c_I\|h\|^2_{H^2(\Omega)}\\ &\quad + \frac{1}{4\epsilon_1}\|\nabla v\|^2_{L_6(\Omega)}\|\operatorname{rot} h\|^2_{L_2(\Omega)}\\ &\leq \epsilon_1 c_{\Omega,I}\|\triangle h\|^2_{L_2(\Omega)} + \frac{1}{4\epsilon_1}\|\nabla v\|^2_{L_6(\Omega)}\|\operatorname{rot} h\|^2_{L_2(\Omega)},\end{aligned}$$

where we used that $\|\nabla h\|_{L_3(\Omega)} \leq \|\nabla h\|^{\frac12}_{L_2(\Omega)}\|\nabla h\|^{\frac12}_{L_6(\Omega)} \leq c_I\|\nabla h\|_{H^1(\Omega)}$. The last inequality above is justified in light of Lemma 4.6.

For the second term on the right-hand side in (5.9) we simply have

$$\|h\|_{L_3(\Omega)}\|\nabla v\|_{L_6(\Omega)}\|\triangle h\|_{L_2(\Omega)} \leq \epsilon_2\|\triangle h\|^2_{L_2(\Omega)} + \frac{1}{4\epsilon_2}\|h\|_{L_2(\Omega)}\|h\|_{L_6(\Omega)}\|\nabla v\|^2_{L_6(\Omega)}.$$

The third term is estimated as follows

$$2\nu_r\int_\Omega \operatorname{rot}\theta\cdot\triangle h\,\mathrm{d}x \leq 2\nu_r\|\operatorname{rot}\theta\|_{L_2(\Omega)}\|\triangle h\|_{L_2(\Omega)} \leq 2\nu_r\epsilon_3\|\triangle h\|^2_{L_2(\Omega)} + \frac{\nu_r}{2\epsilon_3}\|\operatorname{rot}\theta\|^2_{L_2(\Omega)}.$$

Finally, for the fourth term we have

$$\int_\Omega f_{,x_3}\cdot\triangle h\,\mathrm{d}x \leq \|f_{,x_3}\|_{L_2(\Omega)}\|\triangle h\|_{L_2(\Omega)} \leq \epsilon_4\|\triangle h\|^2_{L_2(\Omega)} + \frac{1}{4\epsilon_4}\|f_{,x_3}\|^2_{L_2(\Omega)}.$$

Setting $\epsilon_1 c_{\Omega,I} = \epsilon_2 = \epsilon_4 = \frac{\nu}{6}$ and $\epsilon_3 = \frac12$ yields

$$\begin{aligned}(5.10)\quad \frac12\frac{\mathrm{d}}{\mathrm{dt}}\int_\Omega|\operatorname{rot} h|^2\,\mathrm{d}x + \frac{\nu}{2}\|\triangle h\|^2_{L_2(\Omega)} &\leq \frac{3c_{\Omega,I}}{2\nu}\|\nabla v\|^2_{L_6(\Omega)}\|\operatorname{rot} h\|^2_{L_2(\Omega)}\\ &\quad + \frac{3}{2\nu}\|h\|_{L_2(\Omega)}\|h\|_{L_6(\Omega)}\|\nabla v\|^2_{L_6(\Omega)} + \nu_r\|\operatorname{rot}\theta\|^2_{L_2(\Omega)} + \frac{3}{2\nu}\|f_{,x_3}\|^2_{L_2(\Omega)}\\ &\leq \frac{3c_{I,\Omega}}{\nu}\|\nabla v\|^2_{L_6(\Omega)}\|\operatorname{rot} h\|^2_{L_2(\Omega)} + \nu_r\|\operatorname{rot}\theta\|^2_{L_2(\Omega)} + \frac{3}{2\nu}\|f_{,x_3}\|^2_{L_2(\Omega)},\end{aligned}$$

where in the last inequality we used $\|h\|_{L_2(\Omega)}\|h\|_{L_6(\Omega)} \leq c_I\|h\|^2_{H^1(\Omega)}$ and subsequently utilized Lemma 4.2. Since $\|\operatorname{rot}\theta\|^2_{L_2(\Omega)} \leq 2\|\theta\|^2_{H^1(\Omega)}$ we get from (5.10)

$$\begin{aligned}(5.11)\quad \frac12\frac{\mathrm{d}}{\mathrm{dt}}\int_\Omega|\operatorname{rot} h|^2\,\mathrm{d}x + \frac{\nu}{2}\|\triangle h\|^2_{L_2(\Omega)}&\\ \leq \frac{3c_{\Omega,I}}{\nu}\|\nabla v\|^2_{L_6(\Omega)}\|\operatorname{rot} h\|^2_{L_2(\Omega)} &+ 2\nu_r\|\theta\|^2_{H^1(\Omega)} + \frac{3}{2\nu}\|f_{,x_3}\|^2_{L_2(\Omega)}.\end{aligned}$$

In [Now12, Proof of Lemma 8.4] we established

$$\begin{aligned}\frac12\frac{\mathrm{d}}{\mathrm{dt}}\left(\int_\Omega h^2\,\mathrm{d}x + \int_\Omega\theta^2\,\mathrm{d}x\right) &+ \frac{\nu}{2c_\Omega}\|h\|^2_{H^1(\Omega)} + \frac{\alpha}{2c_\Omega}\|\theta\|^2_{H^1(\Omega)}\\ &\leq \frac{c_{I,\Omega}}{\nu}\|\nabla v\|^2_{L_2(\Omega)}\|h\|^2_{L_3(\Omega)} + \frac{c_{I,\Omega}}{\alpha}\|\nabla\omega\|^2_{L_2(\Omega)}\|h\|^2_{L_3(\Omega)}\\ &\quad + \frac{c_{I,\Omega}}{\nu}\|f_{,x_3}\|^2_{L_{\frac65}(\Omega)} + \frac{c_{I,\Omega}}{\alpha}\|g_{,x_3}\|^2_{L_{\frac65}(\Omega)}.\end{aligned}$$

Multiplying it by $\frac{8\nu_r c_\Omega}{\alpha}$ and adding to (5.11) yields

$$\begin{aligned}\frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}\left(\int_\Omega |\mathrm{rot}\, h|^2 + \frac{8\nu_r c_\Omega}{\alpha}|h|^2 + \frac{8\nu_r c_\Omega}{\alpha}\theta^2\,\mathrm{d}x\right) + \frac{\nu}{2}\|\triangle h\|^2_{L_2(\Omega)} + \frac{4\nu\nu_r c_\Omega}{\alpha}\|h\|^2_{H^1(\Omega)} + 4\nu_r\|\theta\|^2_{H^1(\Omega)}\\
\le \frac{8\nu_r c_{I,\Omega}}{\nu\alpha}\|\nabla v\|^2_{L_2(\Omega)}\|h\|^2_{L_3(\Omega)} + \frac{8\nu_r c_{I,\Omega}}{\alpha^2}\|\nabla\omega\|^2_{L_2(\Omega)}\|h\|^2_{L_3(\Omega)}\\
+ \frac{8\nu_r c_{I,\Omega}}{\nu\alpha}\|f_{,x_3}\|^2_{L_{\frac{6}{5}}(\Omega)} + \frac{8\nu_r c_{I,\Omega}}{\alpha^2}\|g_{,x_3}\|^2_{L_{\frac{6}{5}}(\Omega)}\\
+ \frac{3c_{\Omega,I}}{\nu}\|\nabla v\|^2_{L_6(\Omega)}\|\mathrm{rot}\, h\|^2_{L_2(\Omega)} + 2\nu_r\|\theta\|^2_{H^1(\Omega)} + \frac{3}{2\nu}\|f_{,x_3}\|^2_{L_2(\Omega)}.\end{aligned}$$

Interpolation between L_2 and L_6 shows that $\|h\|_{L_3(\Omega)} \le \|h\|^{\frac{1}{2}}_{L_2(\Omega)}\|h\|^{\frac{1}{2}}_{L_6(\Omega)} \le c_I\|h\|_{H^1(\Omega)}$, because

$$\frac{1}{3} = \frac{\kappa}{2} + \frac{1-\kappa}{6} \quad\Leftrightarrow\quad \kappa = \frac{1}{2}.$$

In view of (4.4) we see that $\|h\|_{H^1(\Omega)} \le c_I\|\mathrm{rot}\, h\|_{L_2(\Omega)}$. Since

$$\begin{aligned}\|f_{,x_3}\|^2_{L_{\frac{6}{5}}(\Omega)} &\le c_\Omega\|f_{,x_3}\|^2_{L_2(\Omega)},\\
\|g_{,x_3}\|^2_{L_{\frac{6}{5}}(\Omega)} &\le c_\Omega\|g_{,x_3}\|^2_{L_2(\Omega)}\end{aligned}$$

we get

$$\begin{aligned}\frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}\left(\int_\Omega |\mathrm{rot}\, h|^2 + \frac{8\nu_r c_\Omega}{\alpha}|h|^2 + \frac{8\nu_r c_\Omega}{\alpha}\theta^2\,\mathrm{d}x\right) + \frac{\nu}{2}\|\triangle h\|^2_{L_2(\Omega)} + \frac{4\nu\nu_r c_\Omega}{\alpha}\|h\|^2_{H^1(\Omega)}\\
+ 2\nu_r\|\theta\|^2_{H^1(\Omega)}\\
\le \left(\frac{8\nu_r c_{I,\Omega}}{\nu\alpha}\|\nabla v\|^2_{L_2(\Omega)} + \frac{8\nu_r c_{I,\Omega}}{\alpha^2}\|\nabla\omega\|^2_{L_2(\Omega)} + \frac{3c_{\Omega,I}}{\nu}\|\nabla v\|^2_{L_6(\Omega)}\right)\|\mathrm{rot}\, h\|^2_{L_2(\Omega)}\\
+ \frac{16\nu_r c_{I,\Omega} + 3\alpha}{2\nu\alpha}\|f_{,x_3}\|^2_{L_2(\Omega)} + \frac{8\nu_r c_{I,\Omega}}{\alpha^2}\|g_{,x_3}\|^2_{L_2(\Omega)}.\end{aligned} \tag{5.12}$$

From (4.4) and natural embeddings it follows

$$\frac{\nu}{2}\|\triangle h\|^2_{L_2(\Omega)} \ge \frac{\nu}{2c_\Omega}\|h\|^2_{H^2(\Omega)} \ge \frac{\nu}{2c_\Omega}\|h\|^2_{H^1(\Omega)} \ge \frac{\nu}{4c_\Omega}\|\mathrm{rot}\, h\|^2_{L_2(\Omega)}.$$

In the same manner

$$\begin{aligned}\frac{4\nu\nu_r c_\Omega}{\alpha}\|h\|^2_{H^1(\Omega)} &\ge \frac{4\nu\nu_r c_\Omega}{\alpha}\|h\|^2_{L_2(\Omega)},\\
2\nu_r\|\theta\|^2_{H^1(\Omega)} &\ge 2\nu_r\|\theta\|^2_{L_2(\Omega)}.\end{aligned}$$

Thus

$$\begin{aligned}\frac{\nu}{4c_\Omega}\|\mathrm{rot}\, h\|^2_{L_2(\Omega)} + \frac{4\nu\nu_r c_\Omega}{\alpha}\|h\|^2_{L_2(\Omega)} + 2\nu_r\|\theta\|^2_{L_2(\Omega)}\\
\ge \frac{\nu}{4c_\Omega}\left(\|\mathrm{rot}\, h\|^2_{L_2(\Omega)} + \frac{16\nu_r c_\Omega^2}{\alpha}\|h\|^2_{L_2(\Omega)} + \frac{8\nu_r c_\Omega}{\nu}\|\theta\|^2_{L_2(\Omega)}\right)\\
= \frac{\nu}{4c_\Omega}\left(\|\mathrm{rot}\, h\|^2_{L_2(\Omega)} + 2c_\Omega\frac{8\nu_r c_\Omega}{\alpha}\|h\|^2_{L_2(\Omega)} + \frac{\alpha}{\nu}\frac{8\nu_r c_\Omega}{\alpha}\|\theta\|^2_{L_2(\Omega)}\right)\\
\ge \frac{\min\{\nu, 2\nu c_\Omega, \alpha\}}{4c_\Omega}\left(\|\mathrm{rot}\, h\|^2_{L_2(\Omega)} + \frac{8\nu_r c_\Omega}{\alpha}\|h\|^2_{L_2(\Omega)} + \frac{8\nu_r c_\Omega}{\alpha}\|\theta\|^2_{L_2(\Omega)}\right).\end{aligned}$$

Denote

$$X(t) = \|\mathrm{rot}\, h(t)\|^2_{L_2(\Omega)} + \frac{8\nu_r c_\Omega}{\alpha}\|h(t)\|^2_{L_2(\Omega)} + \frac{8\nu_r c_\Omega}{\alpha}\|\theta(t)\|^2_{L_2(\Omega)}.$$

Then (5.12) becomes

$$\begin{aligned}\frac{1}{2}\frac{\mathrm{d}}{\mathrm{dt}}X(t) + \frac{\min\{\nu, 2\nu c_\Omega, \alpha\}}{4c_\Omega}X(t)&\\ \leq \left(\frac{8\nu_r c_{I,\Omega}}{\nu\alpha}\|\nabla v\|^2_{L_2(\Omega)} + \frac{8\nu_r c_{I,\Omega}}{\alpha^2}\|\nabla\omega\|^2_{L_2(\Omega)} + \frac{3c_{\Omega,I}}{\nu}\|\nabla v\|^2_{L_6(\Omega)}\right)X(t)&\\ + \frac{16\nu_r c_{I,\Omega}+3\alpha}{2\nu\alpha}\|f_{,x_3}\|^2_{L_2(\Omega)} + \frac{8\nu_r c_{I,\Omega}}{\alpha^2}\|g_{,x_3}\|^2_{L_2(\Omega)}.&\end{aligned}$$

Let

$$\begin{aligned}c_1 &= \frac{\min\{\nu, 2\nu c_\Omega, \alpha\}}{2c_\Omega},\\ c_2 &= \max\left\{\frac{16\nu_r c_{I,\Omega}}{\nu\alpha}, \frac{16\nu_r c_{I,\Omega}}{\alpha^2}, \frac{6c_{\Omega,I}}{\nu}\right\},\\ c_3 &= \max\left\{\frac{16\nu_r c_{I,\Omega}+3\alpha}{\nu\alpha}, \frac{16\nu_r c_{I,\Omega}}{\alpha^2}\right\}.\end{aligned}$$

Then, the last inequality implies that

$$\begin{aligned}\frac{\mathrm{d}}{\mathrm{dt}}\left(X(t)e^{c_1t - c_2\left(\int_{t_0}^t\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}+\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s\right)}\right)&\\ \leq c_3\left(\|f_{,x_3}\|^2_{L_2(\Omega)}+\|g_{,x_3}\|^2_{L_2(\Omega)}\right)e^{c_1t - c_2\left(\int_{t_0}^t\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}+\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s\right)}.&\end{aligned}$$

Integrating with respect to $t \in (t_0, t_1)$ yields

$$\begin{aligned}&X(t_1)e^{c_1t_1 - c_2\left(\int_{t_0}^{t_1}\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}+\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s\right)}\\ &\leq c_3\int_{t_0}^{t_1}\left(\|f_{,x_3}(t)\|^2_{L_2(\Omega)}+\|g_{,x_3}(t)\|^2_{L_2(\Omega)}\right)e^{c_1t - c_2\left(\int_{t_0}^t\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}+\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s\right)}\,\mathrm{d}t\\ &\qquad + X(t_0)e^{c_1t_0}.\end{aligned}$$

Since

$$\int_{t_0}^t\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}+\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s \geq 0$$

we get that

$$\begin{aligned}X(t_1) \leq c_3&e^{-c_1t_1 + c_2\left(\int_{t_0}^{t_1}\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}+\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s\right)}\\ &\cdot\int_{t_0}^{t_1}\left(\|f_{,x_3}(t)\|^2_{L_2(\Omega)}+\|g_{,x_3}(t)\|^2_{L_2(\Omega)}\right)e^{c_1t}\,\mathrm{d}t\\ &+ X(t_0)e^{-c_1(t_1-t_0) + c_2\left(\int_{t_0}^{t_1}\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}+\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s\right)}.\end{aligned}$$

By assumption on $f_{,x_3}$ and $g_{,x_3}$ we obtain

$$\begin{aligned}X(t_1) \leq c_3&e^{-c_1t_1 + c_2\left(\int_{t_0}^{t_1}\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}+\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s\right)}\\ &\cdot\int_{t_0}^{t_1}e^{c_1t}e^{-(t-t_0)}\,\mathrm{d}t\left(\|f_{,x_3}(t_0)\|^2_{L_2(\Omega)}+\|g_{,x_3}(t_0)\|^2_{L_2(\Omega)}\right)\\ &+ X(t_0)e^{-c_1(t_1-t_0) + c_2\left(\int_{t_0}^{t_1}\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}+\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s\right)}.\end{aligned}\tag{5.13}$$

Next we estimate the integral with respect to t

$$\begin{aligned} e^{-c_1t_1}\int_{t_0}^{t_1} e^{c_1t}e^{-(t-t_0)}\,\mathrm{d}t &= e^{-c_1t_1+t_0}\int_{t_0}^{t_1} e^{t(c_1-1)}\,\mathrm{d}t = e^{-c_1t_1+t_0}\frac{1}{c_1-1}e^{t(c_1-1)}\Big|_{t_0}^{t_1} \\ &= e^{-c_1t_1+t_0}\frac{1}{c_1-1}\left(e^{t_1(c_1-1)}-e^{t_0(c_1-1)}\right) \\ &\le \begin{cases} e^{-c_1t_1+t_0}\frac{1}{c_1-1}e^{c_1t_1-t_1} = \frac{1}{c_1-1}e^{-(t_1-t_0)} & \text{for } c_1>1, \\ e^{-c_1t_1+t_0}\frac{1}{1-c_1}e^{c_1t_0-t_0} = \frac{1}{1-c_1}e^{-c_1(t_1-t_0)} & \text{for } c_1<1. \end{cases} \end{aligned}$$

Consider the quantity

$$\int_{t_0}^{t_1}\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}+\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s. \tag{5.14}$$

In view of Lemma 4.7 we see

$$\begin{aligned} \int_{t_0}^{t_1}\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}\,\mathrm{d}s &\le c_{\nu,\alpha,I,\Omega}E^2_{v,\omega}(t_1) \\ &\le \int_{t_0}^{t_1}\|f(s)\|^2_{L_2(\Omega)}+\|g(s)\|^2_{L_2(\Omega)}\,\mathrm{d}s+\|v(t_0)\|^2_{L_2(\Omega)}+\|\omega(t_0)\|^2_{L_2(\Omega)} \\ &\le \int_{t_0}^{t_1}e^{-(s-t_0)}\,\mathrm{d}s\left(\|f(t_0)\|^2_{L_2(\Omega)}+\|g(t_0)\|^2_{L_2(\Omega)}\right)+\|v(t_0)\|^2_{L_2(\Omega)}+\|\omega(t_0)\|^2_{L_2(\Omega)} \\ &\le \|f(t_0)\|^2_{L_2(\Omega)}+\|g(t_0)\|^2_{L_2(\Omega)}+\|v(t_0)\|^2_{L_2(\Omega)}+\|\omega(t_0)\|^2_{L_2(\Omega)}. \end{aligned}$$

To estimate the last term in (5.14) we use the estimate from Theorem 2

$$\begin{aligned} \int_{t_0}^{t_1}\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s &\le c_I\|v\|^2_{L_2(t_0,t_1;H^2(\Omega))}\le c_I\|v\|^2_{W^{2,1}_2(\Omega^{t_1})} \\ &\le c_{\alpha,\nu,\nu_r,I,P,\Omega}\Big(E_{v,\omega}(t_1)+E_{h,\theta}(t_1)+\|f'\|_{L_2(\Omega^{t_1})}+\|v(t_0)\|_{H^1(\Omega)}+1\Big)^6. \end{aligned}$$

By assumption on f, g, their derivative with respect to x_3 we see that

$$\begin{aligned} E_{v,\omega}(t_1)+E_{h,\theta}(t_1)\le c_\Omega\Big(&\|v(t_0)\|_{H^1(\Omega)}+\|\omega(t_0)\|_{H^1(\Omega)}+\|f(t_0)\|_{L_2(\Omega)}+\|g(t_0)\|_{L_2(\Omega)} \\ &+\|f_{,x_3}(t_0)\|_{L_2(\Omega)}+\|g_{,x_3}(t_0)\|_{L_2(\Omega)}\Big). \end{aligned}$$

Thus,

$$\begin{aligned} \int_{t_0}^{t_1}\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s\le\Big(&\|v(t_0)\|_{H^1(\Omega)}+\|\omega(t_0)\|_{H^1(\Omega)}+\|f(t_0)\|_{L_2(\Omega)}+\|g(t_0)\|_{L_2(\Omega)} \\ &+\|f_{,x_3}(t_0)\|_{L_2(\Omega)}+\|g_{,x_3}(t_0)\|_{L_2(\Omega)}+1\Big)^6 \end{aligned}$$

and therefore for t_1 large enough we have

$$-\min\{1,c_1\}(t_1-t_0)+c_2\int_{t_0}^{t_1}\|\nabla v(s)\|^2_{L_2(\Omega)}+\|\nabla\omega(s)\|^2_{L_2(\Omega)}+\|\nabla v(s)\|^2_{L_6(\Omega)}\,\mathrm{d}s<0,$$

which combined with (5.13) implies that

$$X(t_1)\le c_{\alpha,\nu,\nu_r,I,P,\Omega}(t_1)X(t_0),$$

where the function $c_{\alpha,\nu,\nu_r,I,P,\Omega}(t_1)$ has the property

$$\lim_{t_1\to\infty}c_{\alpha,\nu,\nu_r,I,P,\Omega}(t_1)=0.$$

From the above inequality we immediately get that

$$\begin{aligned} \min\left\{1,\frac{8\nu_rc_\Omega}{\alpha}\right\}&\left(\|\mathrm{rot}\,h(t_1)\|^2_{L_2(\Omega)}+\|h(t_1)\|^2_{L_2(\Omega)}+\|\theta(t_1)\|^2_{L_2(\Omega)}\right) \\ &\le c_{\alpha,\nu,\nu_r,I,P,\Omega}\max\left\{1,\frac{8\nu_rc_\Omega}{\alpha}\right\}\left(\|\mathrm{rot}\,h(t_0)\|^2_{L_2(\Omega)}+\|h(t_0)\|^2_{L_2(\Omega)}+\|\theta(t_0)\|^2_{L_2(\Omega)}\right). \end{aligned}$$

For $t_1 \geq t^*$ the inequality yields

$$\begin{aligned}\|\operatorname{rot} h(t_1)\|^2_{L_2(\Omega)} + \|h(t_1)\|^2_{L_2(\Omega)} + \|\theta(t_1)\|^2_{L_2(\Omega)}\\ \leq \|\operatorname{rot} h(t_0)\|^2_{L_2(\Omega)} + \|h(t_0)\|^2_{L_2(\Omega)} + \|\theta(t_0)\|^2_{L_2(\Omega)},\end{aligned}$$

which is our claim. ☐

Proof of Theorem 1. From Lemma 5.4 it follows that

$$\begin{aligned}\sup_k \left(\|\operatorname{rot} h(kT)\|^2_{L_2(\Omega)} + \|h(kT)\|^2_{L_2(\Omega)} + \|\theta(kT)\|^2_{L_2(\Omega)}\right)\\ \leq \|\operatorname{rot} h(0)\|^2_{L_2(\Omega)} + \|h(0)\|^2_{L_2(\Omega)} + \|\theta(0)\|^2_{L_2(\Omega)}.\end{aligned}$$

By assumption

$$\begin{aligned}\sup_k \|f_{,x_3}\|_{L_2(\Omega\times(kT,(k+1)T))} < \infty,\\ \sup_k \|g_{,x_3}\|_{L_2(\Omega\times(kT,(k+1)T))} < \infty.\end{aligned}$$

Thus, we set

$$\sup_k \delta(kT) \leq \delta(0) =: \delta.$$

Let $t_0 = 0$, $t_1 = T$. In view of Theorem 2 with δ we get the existence of regular solution on the interval $[0, T]$. Lemma 5.2 yields the inequality

$$\|v(T)\|^2_{H^1(\Omega)} + \|\omega(T)\|^2_{H^1(\Omega)} < \|v(0)\|^2_{H^1(\Omega)} + \|\omega(0)\|^2_{H^1(\Omega)}$$

for $T > 0$ sufficiently large. It allows us to use Theorem 2 with the initial conditions $v(T)$, $\omega(T)$ and with δ on the time interval $[T, 2T]$. From Lemma 5.2 it follows that

$$\|v(2T)\|^2_{H^1(\Omega)} + \|\omega(2T)\|^2_{H^1(\Omega)} < \|v(T)\|^2_{H^1(\Omega)} + \|\omega(T)\|^2_{H^1(\Omega)},$$

which in view of previous estimate provides us with regular solution on the interval $[0, 2T]$. Reiterating this procedure yields the existence of regular solution on $[0, kT]$. Passing with k up to infinity provides the global existence. ☐

REFERENCES

[ATS73] T. Ariman, M.A. Turk, and N.D. Sylvester, *Microcontinuum fluid mechanics — a review.*, Int. J. Eng. Sci. **11** (1973), 905–930.

[BS70] G.C. Boulougouris and J. Sevilla, *Velocity distribution and other characteristics of steady and pulsatille blood in fine glas tubes.*, Biorheology **7** (1970), no. 2, 85–107.

[CD00] J.W. Cholewa and T. Dłotko, *Global attractors in abstract parabolic problems.*, Cambridge: Cambridge University Press, 2000.

[CMR98] T. Clopeau, A. Mikelić, and R. Robert, *On the vanishing viscosity limit for the 2d incompressible navier-stokes equations with the friction type boundary conditions*, Nonlinearity **11** (1998), no. 6, 1625–1636.

[Eri66] A.C. Eringen, *Theory of micropolar fluids.*, J. Math. Mech. **16** (1966), 1–16.

[Kel06] J.P. Kelliher, *Navier-stokes equations with navier boundary conditions for a bounded domain in the plane.*, SIAM J. Math. Anal. **38** (2006), no. 1, 210–232.

[Lan76] H. Lange, *Die Existenz von Lösungen der Gleichungen, welche die Strömung inkompressibler mikropolarer Flüssigkeiten beschreiben.*, Z. Angew. Math. Mech. **56** (1976), no. 4, 129–139.

[Lan77] ———, *The existence of instationary flows in incompressible micropolar fluids.*, Arch. Mech. (Arch. Mech. Stos.) **29** (1977), no. 5, 741–744.

[LL87] L.D. Landau and E.M. Lifshitz, *Fluid Mechanics: Volume 6 (Course of Theoretical Physics).*, A Butterworth-Heinemann Title; 2 edition, 1987.

[LSU67] O.A. Ladyženskaja, V.A. Solonnikov, and N.N. Ural'ceva, *Linear and quasilinear equations of parabolic type*, Translated from the Russian by S. Smith. Translations of Mathematical Monographs, Vol. 23, American Mathematical Society, Providence, R.I., 1967.

[Łuk89] G. Łukaszewicz, *On the existence, uniquenss and asymptotic properties for solutions of flows of asymmetric fluids.*, Rend. Accad. Naz. Sci. Detta XL, V. Ser. **13** (1989), no. 1, 105–120.

[Łuk99] ———, *Micropolar fluids. theory and applications.*, Boston: Birkhäuser, 1999.

[Mig84] N.P. Migun, *On hydrodynamic boundary conditions for microstructural fluids.*, Rheol. Acta **23** (1984), 575–581.

[Now12] B. Nowakowski, *Large time existence of strong solutions to micropolar equations in cylindrical domains*, ArXiv e-prints (2012), Submitted.

[OTRM97] E.E. Ortega-Torres and M.A. Rojas-Medar, *Magneto-micropolar fluid motion: Global existence of strong solutions.*, Abstr. Appl. Anal. **4** (1997), no. 2, 109–125.

[PRU74] A.S. Popel, S.A Regirer, and P.I. Usick, *A continuum model of blood flow.*, Biorheology **11** (1974), 427–437.

[RM97] M.A. Rojas-Medar, *Magneto-micropolar fluid motion: Existence and uniqueness of strong solution.*, Math. Nachr. **188** (1997), 301–319.

[RZ08] J. Rencławowicz and W.M. Zajączkowski, *Large time regular solutions to the navier-stokes equations in cylindrical domains.*, Topol. Methods Nonlinear Anal. **32** (2008), no. 1, 69–87.

[SASM02] K.V. Sharp, R.J. Adrian, J.G. Santiago, and J.I. Molho, *Liquid Flows in Microchannels.*, The MEMS Handbook (Mohamed Gad ed Hak, ed.), CRC Press, 2002.

[Sav78] V.A. Sava, *The initial-boundary-value problems in the theory of micropolar fluids.*, Z. Angew. Math. Mech. **58** (1978), 511–518.

[Sol65] V.A. Solonnikov, *On boundary value problems for linear parabolic systems of differential equations of general form.*, Proc. Steklov Inst. Math. **83** (1965), 1–184, Translation from: Trudy Mat. Inst. Steklov 83 (1965), 3–163, (Russian).

[Sol73] ———, *Overdetermined elliptic boundary-value problems.*, Journal of Mathematical Sciences **1** (1973), no. 4, 477–512, Translation from: Zap. Nauchn. Sem. LOMI 21 (1971), 112–158, (Russian).

[Tem79] R. Temam, *Navier-Stokes equations.*, revised ed., Studies in Mathematics and its Applications, vol. 2, North-Holland Publishing Co., Amsterdam, 1979, Theory and numerical analysis, With an appendix by F. Thomasset.

[Yam05] N. Yamaguchi, *Existence of global strong solution to the micropolar fluid system in a bounded domain.*, Math. Methods Appl. Sci. **28** (2005), no. 13, 1507–1526.

[Zaj05] W.M. Zajączkowski, *Long time existence of regular solutions to navier-stokes equations in cylindrical domains under boundary slip conditions.*, Stud. Math. **169** (2005), no. 3, 243–285.

[Zaj11] ———, *On global regular solutions to the Navier-Stokes equations in cylindrical domains.*, Topol. Methods Nonlinear Anal. **37** (2011), no. 1, 55–86.

Bernard Nowakowski, Institute of Mathematics, Polish Academy of Sciences, Śniadeckich 8, 00-956 Warsaw, Poland
E-mail address: `bernard@impan.pl`